



## Fertilization trajectory of the root crop *Raphanus sativus* across atmospheric $p\text{CO}_2$ estimates of the next 300 years

Brian A. Schubert\*, A. Hope Jahren

Department of Geology and Geophysics, University of Hawaii, 1680 East-West Road, POST 701, Honolulu, HI 96822, USA

### ARTICLE INFO

#### Article history:

Received 28 September 2010

Received in revised form

24 November 2010

Accepted 25 November 2010

Available online 17 December 2010

#### Keywords:

Biomass  
Elevated carbon dioxide  
Radish  
Sustainability

### ABSTRACT

Plant biomass is known to increase in response to elevated atmospheric  $\text{CO}_2$  concentration ( $p\text{CO}_2$ ); however, no experiments have quantified the trajectory of crop fertilization across the full range of  $p\text{CO}_2$  levels estimated for the next 300 years. Here we quantify the above- and below-ground biomass response of *Raphanus sativus* (common radish) across eight  $p\text{CO}_2$  levels ranging from 348 to 1791 ppmv. We observed a large net biomass increase of 58% above ground and 279% below ground. A large part of the net increase (38% of the above-ground and 53% of the below-ground) represented biomass fertilization at the high levels of  $p\text{CO}_2$  (700–1791 ppmv) predicted if fossil fuel emissions continue unabated. The trajectory of below-ground fertilization in *R. sativus* greatly exceeded a trajectory based on extrapolation of previous experiments for plants grown at  $p\text{CO}_2 < 800$  ppmv. Based on the experimental parameters used to grow these plants, we hypothesize that these experiments represent the maximum  $\text{CO}_2$  fertilization that can be achieved for this plant growing under low light levels. If the below-ground biomass enhancement that we have quantified for *R. sativus* represents a generalized root-crop response that can be extrapolated to agricultural systems, below-ground fertilization under very high  $p\text{CO}_2$  levels could dramatically augment crop production in some of the poorest nations of the world, provided that water resources are sufficient and sustainable.

© 2010 Elsevier B.V. All rights reserved.

### 1. Introduction

The effects of climate change on agroecosystem sustainability and crop production are difficult to predict (Fuhrer, 2003), but have great implications for the global food supply (Schmidhuber and Tubiello, 2007). Tradeoffs between increased plant-tissue production (i.e., “ $\text{CO}_2$  fertilization”, Lobell and Field, 2008) and increased irrigation requirements resulting from climate change make the issue complex (Melillo et al., 1993; Tubiello et al., 2007). Current understanding of  $\text{CO}_2$  fertilization is based on thousands of two-point experiments that have grown hundreds of plant species under ambient (300–420 ppmv) vs. slightly elevated (475–750 ppmv)  $p\text{CO}_2$ . Taken together, these experiments revealed an average total biomass increase of 20–54% (Kimball et al., 1993; Poorter, 1993; Wand et al., 1999; Poorter and Navas, 2003; Ainsworth and Long, 2005). The elevated  $p\text{CO}_2$  levels used for these plant-growth experiments generally ranged from 475 to 750 ppmv, and were based on early projections for the year 2100 (Houghton et al., 1992); recent projections include  $p\text{CO}_2$  estimates >750 ppmv (e.g., 560–820 ppmv for 2100, Nakicenovic and Swart, 2000). By the year 2300,  $p\text{CO}_2$  levels may reach 1800 ppmv

if fossil fuel emissions continue unabated (Zachos et al., 2008). A simple extrapolation of  $p\text{CO}_2$  data from Mauna Loa (1959–2008) NOAA/ESRL (2010), fitted to a second order polynomial equation ( $R^2 = 0.999$ ), projects  $p\text{CO}_2 = 1847$  ppmv for the year 2300. Although some two-point experiments have compared biomass in plants grown under ambient conditions to very high  $p\text{CO}_2$  levels (i.e., 900–1200 ppmv) (Tognoni et al., 1966; Knecht, 1975; Luxmoore et al., 1986; Stuhlfauth et al., 1987; Riechers and Strain, 1988; Wheeler and Tibbitts, 1989; Vessey et al., 1990; Lord et al., 1993) and  $p\text{CO}_2$  levels beyond those that plants have ever experienced on Earth (i.e., >10,000 ppm) (Park and Epstein, 1960; Schwarz and Strain, 1990; R.M. Wheeler et al., 1994), fertilization has not been quantified for any plant grown under the same conditions at multiple levels of  $p\text{CO}_2$  rise approaching the full range of greenhouse gas estimates for the next 300 years. The  $\text{CO}_2$  emissions of the world’s most populated countries are in an extreme state of flux (United Nations MDGI, 2010), which confers a large uncertainty to estimates for rising  $p\text{CO}_2$  levels. For this reason, the true shape of the relationship between  $p\text{CO}_2$  and plant biomass (i.e., the trajectory) must be known in order to accurately predict effects on crop production under multiple scenarios of  $p\text{CO}_2$ -rise.

The need for accurate root-crop fertilization trajectories is pressing: although production of root crops globally has increased more than 50% in the last 50 years (Godfray et al., 2010), the people most dependent on root-crop agriculture reside in the most improv-

\* Corresponding author. Tel.: +1 808 956 0457; fax: +1 808 956 5512.  
E-mail address: [bschube@hawaii.edu](mailto:bschube@hawaii.edu) (B.A. Schubert).


erished portions of the globe (Kates and Dasgupta, 2007; United Nations FAO, 2010). Ethiopia, one of the ten poorest and most malnourished nations of the world (United Nations Development Programme, 2009), relies on “roots and tubers” as the number-one commodity within its agriculture sector, which generates more than 50% of Ethiopia’s gross domestic product (United Nations FAO). *Raphanus sativus* L. (Cherry Belle radish), in particular, is a crop of some importance. Each year, Americans spend sixty million dollars on grocery-store radishes, and the U.S. earns about 10 million dollars on its radish exports (Lucier and Jerardo, 2004). One-third of one’s daily-recommended allowance of vitamin C can be gained from 20 calories of radish, at a cost of just eleven cents to the U.S. consumer (Otten et al., 2006). Radishes have the potential to be a highly sustainable crop: they grow to marketable size in three weeks, can be irrigated with wastewater (Bakhsh and Hassan, 2005), exist as several disease-resistant wild-types (Snow and Campbell, 2005), and *Raphanus* seed-oil has been proposed as a biofuel (Domingos et al., 2008).

Here we report the above- and below-ground biomass response and carbon stable isotope values for 132 *R. sativus* plants grown across eight carbon dioxide levels ranging from 348 to 1791 ppmv (Table S1) using established growth-chamber techniques (Jahren et al., 2008). We chose the herbaceous plant *R. sativus* based on its long-standing use as a model plant for environmental experiments (Kostka-Rick and Manning, 1993) and its classification in the Brassicaceae family with *Arabidopsis*, which has served as a model for examinations of the genetic basis of responses to elevated  $p\text{CO}_2$  (Li et al., 2008). The response of *Raphanus* spp. to estimates of continued  $p\text{CO}_2$  rise may serve as a harbinger for root crops as we contrast the 21st-century results of Global Change with its 20th-century predictions.

## 2. Materials and methods

### 2.1. Growth conditions

*R. sativus* (Harris Seeds of Rochester, ID # 0071) were grown within positive-pressure Plexiglass chambers under controlled  $p\text{CO}_2$ , light, and temperature after the design described within (Jahren et al., 2008). Carbon dioxide levels were increased by bleeding pure cylinder  $\text{CO}_2$  into the intake pipe where it mixed with the ambient air before reaching the growth chamber. For each chamber the  $\text{CO}_2$  flow rate was precisely controlled with an inline needle valve, thus enabling enrichment of air to the desired concentration of  $\text{CO}_2$ . Air was exhausted through an upper pipe and vented through a fume hood. Complete atmospheric turnover occurred approximately once every 10.4 min. The eight  $p\text{CO}_2$  levels (348, 388, 413, 426, 760, 1090, 1425, and 1791 ppmv) were maintained within 10% of the desired  $p\text{CO}_2$  level during four weeks of growth (Fig. 1A). We grew sixteen to eighteen plants at each  $p\text{CO}_2$  level, which is double the median number of plants harvested in 350 previous elevated- $\text{CO}_2$  experiments (Poorter and Navas, 2003). Experiments consisted of five days of germination on damp paper in the dark under ambient  $p\text{CO}_2$ , followed by four weeks of growth in standard potting soil (Miracle-Gro Moisture Control). This timing was designed to capture maximum growth after the cotyledons ceased to function but before reproduction was initiated, to assess biomass at a commensurate developmental stage of harvest within previous experiments (Poorter and Navas, 2003). Fluorescent grow lamps (33-W GE Brightstik) on adjustable shelving provided up to  $230 \mu\text{mol m}^{-2} \text{s}^{-1}$  photosynthetic photon flux (400–700 nm), as measured at the leaf surface. Lights were controlled by a diurnal timer (11.75 h on; 12.25 h off per 24-h cycle). Because the lights generated heat inside the chambers, higher light levels could not have been achieved without increasing the growing temperature



**Fig. 1.**  $p\text{CO}_2$  (A), temperature (B), and relative humidity (C) conditions for three ambient (black) and three elevated (gray)  $p\text{CO}_2$  experiments. Within the growth chambers,  $p\text{CO}_2$  and temperature conditions were controlled; relative humidity fluctuated naturally. The mean and standard deviation for all experiments are presented in Table S1.

inside the chambers above the temperatures for maximum growth of *R. sativus* (Fig. 2 within Idso and Kimball, 1989). Mean daytime growing temperatures in the chambers averaged  $23.3^\circ\text{C}$  or  $29.5^\circ\text{C}$  (Fig. 1B) and matched temperatures shown for maximum growth of *R. sativus* (Fig. 2 within Idso and Kimball, 1989). Relative humidity was not controlled (average relative humidity was 29% and 37% for the high and low temperature regimes, respectively) (Fig. 1C); therefore, the affect of relative humidity on biomass was not determined. Wong (1993), however, found no significant difference in biomass between *R. sativus* plants grown at low (35%) and high (90%) relative humidity for ambient (350 ppmv) and slightly elevated (700 ppmv)  $p\text{CO}_2$ . Each plant occupied a separate soil container and was supplied with excess water, nitrogen, phosphorous, and trace nutrients. Soil was maintained at “field moist” conditions (Soil Survey Staff and Soil Conservation Service, 1999).

Plant growth within chambers has been documented to yield larger biomass relative to field growth (Baldy, 1986; Long et al., 2006); however, growth chamber design (Leadley and Drake, 1993; Edwards et al., 2009), computational methods (Poorter and Garnier, 1996), and experimental parameters such as light levels, temperature, relative humidity, water, nutrients, and pot size can effect plant growth results (Wong, 1979, 1993; Sionit et al., 1982; Sionit, 1983; Morison and Gifford, 1984; Zangerl and Bazzaz, 1984; Idso and Kimball, 1989; Bhattacharya et al., 1990; Farrar and Williams, 1991; Rogers et al., 1994; T.R. Wheeler et al., 1994; Schenk et al., 1996; Jablonski, 1997). Although our experiments were designed to maximize biomass at each  $p\text{CO}_2$  level (optimal temperature, water,

**Table 1**  
Growth parameters that can affect the CO<sub>2</sub> fertilization response.

Parameter	This study	CO <sub>2</sub> fertilization response <sup>a</sup>
Chamber vs. field	Chamber	+
Light	230 μmol m <sup>-2</sup> s <sup>-1</sup> PFF (400–700 nm)	–
Temperature	23.3 and 29.5 °C (optimal <sup>b</sup> )	+
Relative humidity	29 and 37%	x <sup>c</sup>
Water	Field moist <sup>d</sup>	+
Nutrients	Excess N, P, and trace elements	+
Pot size	0.3 L	–

<sup>a</sup> Maximum (+), limited (–), and no effect (x).

<sup>b</sup> Fig. 2 within *Idso and Kimball (1989)*.

<sup>c</sup> *Wong (1993)*.

<sup>d</sup> *Soil Survey Staff and Soil Conservation Service (1999)*.

and nutrients for growth), it is possible that low light levels and pot artifacts lessened the CO<sub>2</sub> fertilization effect, especially at very high pCO<sub>2</sub> levels, by limiting the maximum, potential biomass (Table 1). Conversely, biomass enhancement of crops grown in agricultural fields is not limited by light levels or pot size, but may be limited by temperature, water, and nutrient stresses. Implications of resource limitations on CO<sub>2</sub> fertilization are discussed below.

## 2.2. Biomass assessment

After four weeks of chamber growth, above-ground tissues (stems and leaves) were harvested and separated from below-ground tissues (hypocotyl and roots), dried for 5 days at 60 °C, and weighed (uncertainty ±1 mg). We report all biomass weights in units of ln-transformed milligrams or as back-transformed values in units of milligrams (Table S3), as is the standard method for comparing plant mean dry weight (*Poorter and Garnier, 1996*).

We analyzed our biomass data in terms of a hyperbolic response, as routinely invoked for modeling CO<sub>2</sub> and light responses of photosynthesis in single leaves and whole plants (*Lloyd and Farquhar, 1996; Nowak et al., 2004*). In order to perform a direct and specific quantitative comparison to lower pCO<sub>2</sub> experiments, the data were plotted and fitted to a two-parameter rectangular hyperbola, after the method used by *Hunt et al. (1991, 1993)* for assessing the trajectory of the biomass response for 36 herbaceous species grown under pCO<sub>2</sub> levels ranging from 365 to 812 ppmv:

$$y = \frac{abx}{a + bx} \quad (1)$$

where  $y$  is the plant-yield variable (i.e., biomass);  $x$  is the pCO<sub>2</sub> level;  $a$  is the asymptote; and  $b$  is the slope at  $y = x = 100$  ppmv, the value used for CO<sub>2</sub> compensation point of dicotyledons (*Ludlow and Jarvis, 1971*; after *Hunt et al., 1991*). *Hunt et al. (1991, 1993)* defined the value  $b$  as an overall index of CO<sub>2</sub> responsiveness: at low  $b$  values the function approaches the asymptote slowly, at high  $b$  more rapidly. Because the function is non-linear, fitted estimates of  $a$  and  $b$  were obtained by iterative optimization to maximize the correlation coefficient ( $r$ ):

$$r = \sqrt{1 - \frac{\sum_{i=1}^n (y_i - y_{f_i})^2}{\sum_{i=1}^n (y_i - \bar{y})^2}} \quad (2)$$

where  $y_i$  is the average biomass value at each pCO<sub>2</sub> level,  $y_{f_i}$  is the fitted biomass value at each pCO<sub>2</sub> level, and  $y_{avg}$  is the mean biomass value for all pCO<sub>2</sub> levels.

**Table 2**  
Values for  $a$ ,  $b$ , and  $r$  (Eqs. (1) and (2), see text) calculated for *Hunt et al. (1991, 1993)* and this study.

	$a$	$b$	$r$
<i>Hunt et al. (1991, 1993)</i>			
Above <sup>a</sup>	6.97	0.38	– <sup>b</sup>
Below <sup>c</sup>	6.40	0.38	–
Total <sup>a</sup>	7.41	0.46	–
This study <sup>d</sup>			
Above	5.89	0.23	0.88
Below	6.18	0.07	0.82
Total	6.68	0.16	0.86

<sup>a</sup> Calculated as the average  $a$  and  $b$  values reported in *Hunt et al. (1991, 1993)* for all responsive species (defined as having  $b < 1$  for total biomass).

<sup>b</sup> Average biomass measurements for each species and pCO<sub>2</sub> level were not reported within *Hunt et al. (1991, 1993)*; therefore,  $r$ -values could not be calculated.

<sup>c</sup> *Hunt et al. (1991, 1993)* did not report  $a$  and  $b$  values for below-ground biomass. We calculated the  $a$  and  $b$  values for below-ground biomass as the difference between the total- and above-ground biomass; total- and above-ground biomass were calculated for each pCO<sub>2</sub> level using Eq. (1) (see text). Fitted estimates of  $a$  and  $b$  were then obtained using Eq. (2), with  $r = 1$ .

<sup>d</sup> Using Eq. (2) (see text) fitted estimates of  $a$  and  $b$  were obtained by iterative optimization to maximize the correlation coefficient  $r$ .

## 2.3. Stable isotope analyses

After biomass assessment, above- and below-ground tissues were ground to a uniform powder in preparation for stable isotope analyses. The δ<sup>13</sup>C values of plant tissues were analyzed using a Eurovector automated combustion system in conjunction with an Isoprime stable isotope mass spectrometer. All samples were introduced to the combustion system in pure tin capsules; stable isotope values are reported in standard δ-notation (‰). The reporting standard is Vienna Pee Dee Formation limestone (VPDB) with  $R = {}^{13}\text{C}/{}^{12}\text{C} = 0.011224$ . Analytical uncertainty associated with each measurement was ±0.1‰. Presented values represent the average of three replicate capsules; the average isotopic variability of replicate capsules was ±0.15‰ for all tissues. Combustion also resulted in a quantification of percent carbon content in each sample.


We sampled gaseous CO<sub>2</sub> in each chamber using a pre-evacuated, 250-cm<sup>3</sup>, glass vessel. We placed the vessel within the chamber we wished to sample, opened the stopcock, and let it equilibrate with the surrounding air for 10 min. We then concentrated and purified the CO<sub>2</sub> from the sampled atmosphere by slowly bleeding the air sample through a vacuum manifold, freezing any water into an ethanol/liquid-nitrogen cooled trap, and then freezing CO<sub>2</sub> into a series of liquid-nitrogen cooled traps, while simultaneously pumping away noncondensable gases. Pure CO<sub>2</sub> sample gas was analyzed for the δ<sup>13</sup>CO<sub>2</sub> value through the dual inlet of the mass spectrometer. We further purified one sample with 0.5 g of reduced copper at 450 °C for 2 h in order to remove N<sub>2</sub>O and thus test the effect of naturally occurring N<sub>2</sub>O on chamber δ<sup>13</sup>CO<sub>2</sub>. We found no difference between the δ<sup>13</sup>CO<sub>2</sub> value of the sample reacted with reduced copper and the non-reacted sample.

## 3. Results

### 3.1. Increases in biomass

In keeping with the majority of the previous studies on CO<sub>2</sub> fertilization, we observed *R. sativus* grown to maturity under elevated pCO<sub>2</sub> levels to have greater total biomass than those grown under ambient atmospheric conditions (Fig. 2A). Using the hyperbolic model described in Section 2.2, we found that while the above-ground response ( $b = 0.23$ ) was similar to the mean value of responsive (i.e., total biomass,  $b < 1$ ) angiosperms ( $b = 0.38$ ;  $n = 23$ )





**Fig. 2.** Percent-biomass-increase observed for *R. sativus* across the full range of atmospheric  $p\text{CO}_2$  levels estimated for the next 300 years (348–1791 ppmv) compared with previous results on root crops (a, *Daucus carota*, carrot; b, *Raphanus sativus*, radish; c, *Beta vulgaris*, sugar beet; d, *Ipomoea batatas*, sweet potato; e, *Solanum tuberosum*, potato) and herbaceous species (Hunt et al., 1991, 1993) at lower  $p\text{CO}_2$  levels. Percent-biomass-increase curves for this study (open diamonds) were calculated from the fitted hyperbolic functions for biomass (Eqs. (1) and (2) within text) using an ambient  $p\text{CO}_2$  level of 389 ppmv (present  $p\text{CO}_2$  level). Percent-biomass-increase curves (closed diamonds) were calculated using Eq. (1) and the mean  $a$  and  $b$  values reported for 23 responsive ( $b < 1$ , total biomass) herbaceous species (Hunt et al., 1991, 1993). Open diamonds (this study) and closed diamonds (Hunt et al., 1991, 1993) mark  $p\text{CO}_2$  levels under which the plants were grown; dashed lines extrapolate data from Hunt et al. (1991, 1993). Root crop data (shaded bars) were compiled from published data on dry biomass following the methods of Poorter (1993). Data were restricted as follows: (1) only data from the vegetative phase were used, (2) only experiments with control  $p\text{CO}_2$ -levels between 300 and 450 ppmv and elevated  $p\text{CO}_2$ -levels less than 2000 ppmv were used, in order to make a direct, quantitative comparison with data from this study, (3) for studies in which multiple environmental variables were tested (i.e., ozone, water availability, irradiance, etc.), biomass data were collected from experiments in which control plants grew best, and (4) for studies with multiple harvest dates, only data from the final harvest were used. These criteria resulted in a total of 127 observations of percent biomass increase ( $n = 38$ , above ground;  $n = 45$ , below ground;  $n = 44$ , total biomass) (see Table S2). Data for root crops extend the range of elevated  $p\text{CO}_2$  levels tested and mark the average biomass increase, relative to ambient  $p\text{CO}_2$  levels. (A) Total biomass (sugar beet,  $n = 14$ ; carrot,  $n = 3$ ; sweet potato,  $n = 5$ ; radish,  $n = 10$ ; potato,  $n = 12$ ). Except for potato (e), total percent biomass increases greater for root crops (shaded bars) than non-crop species (closed diamonds). Note that the average, percent-total-biomass-increase at lower  $p\text{CO}_2$  levels for *Raphanus* from previous studies (shaded bar, b) is similar to that observed in this study (open diamonds), but the percent increase in total biomass at very high  $p\text{CO}_2$  levels observed in this study exceeds the extrapolated data of Hunt et al. (1991, 1993) (dashed curve). (B) The above-ground response for root crops (sugar beet,  $n = 10$ ; sweet potato,  $n = 8$ ; radish,  $n = 6$ ; potato,  $n = 14$ ) does not exceed the response of non-root species at any  $p\text{CO}_2$  level. (C) Below-ground biomass fertilization for root crops (sugar beet,  $n = 10$ ; carrot,  $n = 2$ ; sweet potato,  $n = 5$ ; radish,  $n = 6$ ; potato,  $n = 22$ ) at low  $p\text{CO}_2$  levels (<800 ppmv) (except potato) greatly exceeds the average response of non-crops across the full range of  $p\text{CO}_2$  levels; however, the difference between root and

(Hunt et al., 1991, 1993), the below-ground response was remarkable ( $b = 0.07$ ) (Table 2).

Individual variability is inherent to all biological experiments; however the plant-to-plant biomass variability did not exceed what is commonly observed for plant experiments in general. Total-biomass variability for each set of growing conditions (measured as the standard deviation of the ln-transformed biomass data) averaged 0.29, which is nearly identical to the median variability in total biomass determined by Poorter and Navas (2003) for 700 herbaceous plants (0.28).


Percent biomass increase across the full spectrum of  $p\text{CO}_2$  levels, calculated from the fitted biomass values, showed that the total biomass increase we observed for *R. sativus* at  $p\text{CO}_2$  levels up to 760 ppmv was consistent with the average percent increase observed in ten experiments on *R. sativus* grown at ambient (330–386 ppmv) and elevated (600–761 ppmv)  $p\text{CO}_2$  (Fig. 2A). Maximum biomass augmentation, however, occurred in below-ground tissues, which will likely have a major impact on the distribution of biomass within the plant. An increase in root-to-shoot ratio is a common observation in experiments that elevate  $p\text{CO}_2$  to ~700 ppmv (Rogers et al., 1994; Rogers et al., 1996) resulting from increased root size (length, diameter, and dry weight) (Rogers et al., 1992), and has been observed in *R. sativus*, specifically (Overdieck et al., 1988; Barnes and Pfirrmann, 1992; Chu et al., 1992; Wong, 1993; Jablonski, 1997). The percent biomass increase was up to 58% and 279% for above- and below-ground tissues, respectively (Fig. 2B and C). Much of the net increase (38% for above- and 53% for below-ground biomass) occurred above 700 ppmv, i.e., above the average  $p\text{CO}_2$  estimates for 2100 (Houghton et al., 1992) and most commonly used elevated  $p\text{CO}_2$  level in two-point experiments (Poorter and Navas, 2003).

For above-ground tissues, the trajectory of percent biomass increase observed in our experiments up to ~1800 ppmv is identical to the trajectory of Hunt et al. (1991, 1993) for lower  $p\text{CO}_2$  experiments (<800 ppm) (Fig. 2B). The trajectory that we observed for below-ground biomass increase was extremely high compared to other herbs (Fig. 2C), causing the trajectory for total biomass to similarly exceed extrapolation based on previous experiments (Fig. 2A). If we compare the fertilization trajectory we observed to the results of two-point studies on root crops, we see that the data for lower  $p\text{CO}_2$  studies is within the range of our study for both above- and below-ground tissues (Fig. 2), and for very high  $p\text{CO}_2$  levels, below-ground fertilization increases significantly.

### 3.2. Stable carbon isotope confirmation of growing conditions

Experiments were conducted under luxury conditions (i.e. excess water and nutrients) and temperature levels for optimal growth (Fig. 2 within Idso and Kimball, 1989; Wong, 1993) to allow us to measure maximum biomass enhancement. Nevertheless, it is important to show that lower biomass plants were not somehow less able to access water and nutrients. Multiple studies have shown that water, nitrogen, temperature, light, and other environmental stresses alter the net carbon isotopic fractionation ( $\Delta$ ) between atmospheric  $\delta^{13}\text{C}_2$  and the resultant  $\delta^{13}\text{C}$  value of plant-tissue (Ehleringer et al., 1986, 2002; Ehleringer and Cooper, 1988; Farquhar et al., 1989; Zimmerman and Ehleringer, 1990; Arens et al., 2000; Bowling et al., 2002; Yin et al., 2005; Aranda et al., 2007; Cernusak et al., 2009; Yin et al., 2009; Kranabetter et al., 2010). If environmental stress exerted an influence over biomass production

**Fig. 2.** non-root species is greatest at very high  $p\text{CO}_2$  levels (>800 ppmv). In general, previous data for above-ground, below-ground, and total biomass for *Raphanus* grown under slightly elevated  $p\text{CO}_2$  levels (shaded bars, b) are consistent with data from this study (open diamonds).



**Fig. 3.** Carbon isotope composition of plant tissue compared to plant biomass and to the carbon isotope composition of chamber  $\text{CO}_2$ . Lack of correlation between (A) the net carbon isotope fractionation ( $\Delta = \delta^{13}\text{C}_{\text{plant}} - \delta^{13}\text{CO}_2$ ) and the amount of biomass produced contrasted with the strong correlation between (B) the carbon isotope composition of the plant tissue ( $\delta^{13}\text{C}_{\text{plant}}$ ) and the carbon isotope composition of chamber  $\text{CO}_2$  ( $\delta^{13}\text{CO}_2$ ). Carbon isotope values are given in units of per mil (‰) relative to VPDB standard. Biomass is given as dry weight in units of ln-transformed milligrams [ln(mg)] after Hunt et al. (1991, 1993) and Poorter and Garnier (1996).

in our experiments, we would expect to see a relationship between the mass of tissue produced and the net carbon isotope fractionation between the plant and the atmosphere under which it grew. To the contrary, we saw no correlation (i.e.,  $R^2 < 0.1$ ; Fig. 3A). Elevated  $\text{CO}_2$  may increase water-use efficiency (Eamus, 1991; Lawlor and Mitchell, 1991; Ceulemans and Mousseau, 1994; Jackson et al., 1994; Medlyn et al., 2001), but no isotopic evidence of water or nutrient stress or limitation was found in small or large plants; similarly, evidence for self-shading by leaves was not found. There was a strong correlation, however, between the carbon isotope composition of the plant tissue and the atmosphere under which it grew (i.e.,  $R^2 = 0.99$ ; Fig. 3B). This relationship was expected and is consistent with what we have shown for lower  $p\text{CO}_2$  levels using similarly depleted cylinder- $\text{CO}_2$  to elevate the  $\text{CO}_2$  levels within the chambers (Arens et al., 2000; Jahren et al., 2008).

## 4. Discussion

### 4.1. Implications for agriculture and sustainability

Other workers have shown that below-ground tissues for root crops may increase significantly more than non-root crops under slightly elevated  $p\text{CO}_2$  levels (Idso et al., 1988) because the enlarged

hypocotyl provides a carbon sink (Overdieck et al., 1988; Miglietta et al., 2000); our work suggests that this effect will continue to augment at very high  $p\text{CO}_2$  levels ( $>750$  ppmv). Biomass enhancement due to elevated  $p\text{CO}_2$  has been hypothesized to be ultimately subject to secondary limitation by nutrients (Ceulemans et al., 1999), in particular nitrogen (Finzi et al., 2007). However, increased allocation to below-ground tissues may enhance a plant's ability to acquire nitrogen, water, phosphorous and other micronutrients (Langley and Megonigal, 2010). Researchers have observed a decrease in protein concentration (Taub et al., 2008) and higher C:N ratios in plant tissues grown under elevated  $p\text{CO}_2$  compared to tissues grown under ambient conditions due to increased nitrogen use efficiency at both low and high N supply (Overdieck et al., 1988; Conroy, 1992; Drake et al., 1997; Cotrufo et al., 1998; Taub and Wang, 2008). On average, however, N concentrations may decrease less in below-ground than above-ground tissues (Cotrufo et al., 1998). Other researchers have indicated that nutritive quality of crops is not expected to diminish due to rising  $p\text{CO}_2$  levels (Idso and Idso, 2001). In our experiments, %C in tissues ( $34.7 \pm 2.7\%$  for above-ground tissue;  $36.3 \pm 2.3\%$  for below-ground tissue) was not correlated with  $p\text{CO}_2$  growth-level, in agreement with previous studies (Overdieck and Reining, 1986; Overdieck et al., 1988); therefore, the biomass increase we observed did not carry a proportional increase in N-demand. Root growth also promotes mycorrhizal fungi that enhance availability and uptake of nutrients in the root zone (Lin et al., 2000; van Hees et al., 2006; Javaid, 2009), including in agricultural settings (Zhang et al., 2010). *R. sativus* (Brassicaceae) is part of the minority 8% of plant families that have no symbiotic association with mycorrhizal fungi (Wang and Qiu, 2006); however, root exudates of Brassicaceae plants, including *R. sativus* specifically, have been shown to stimulate hyphal growth of mycorrhizal fungi (Zeng et al., 2003). Because fertilizer is a major and limiting agricultural expense within developing nations FAO (2008), increases in crop yield that do not require proportional fertilizer additions are a critical requirement for global sustainability.

The potential effect of increased below-ground biomass upon irrigation requirements is complex. It is reasonable to expect that any biomass enhancement will require an increase in plant-available water based on long-standing observations of the amount of water required for crop growth (e.g., 200–800 g of  $\text{H}_2\text{O}$  used per production of 1 g of dry tissue, Lyon et al., 1952). However, water-use efficiency in plants increases under elevated  $p\text{CO}_2$  due to a decrease in stomatal conductance (Eamus, 1991; Lawlor and Mitchell, 1991; Chu et al., 1992; Ceulemans and Mousseau, 1994; Jackson et al., 1994; Medlyn et al., 2001). In addition, plant roots increase mineral surface area below ground by breaking, fracturing, and aligning mineral grains (April and Keller, 1990), leading to increased residence time of soil water (Drever, 1994) and possibly buffering plants against droughts (Norby et al., 2004).

## 5. Conclusions

Our experiments showed that under slightly elevated  $p\text{CO}_2$  ( $<760$  ppmv), *R. sativus* biomass increased similarly to previous studies on crop and non-crop species. However, at higher  $p\text{CO}_2$  levels, which include recent estimates for the year 2300 (Zachos et al., 2008), biomass enhancement in the primary, edible organ was much larger than what might be predicted from trajectories of non-crop species at lower- $p\text{CO}_2$  levels; the lower- $p\text{CO}_2$  portion of the trajectory we observed was consistent with two-point studies on root crops. If the dramatic biomass response to very high  $p\text{CO}_2$  levels we have observed for *R. sativus* is shown to exist across other root crops (Miglietta et al., 2000) grown in the field, it may have implications for food production and future greenhouse gas emissions under the most current estimates of  $p\text{CO}_2$

rise for the next 300 years. Yield increases of 135% in the last 50 years, due to agricultural intensification, have avoided emissions of 161 GtC, or 34% of the total carbon emitted by humans between 1850 and 2005 (Burney et al., 2010); further yield increase would continue this trend. Below-ground nutrient limitation could be inherently self-alleviating at elevated  $p\text{CO}_2$  due to positive feedbacks between biomass production and water, nitrogen, and cation availability. The implications for global sustainability are complex: the top-twenty producing countries in the world sell  $7.8 \times 10^9$  kg of roots and tubers each year (potatoes are excluded from this figure). Within these top-producers are four of the world's poorest nations, Ethiopia, Eritrea, the Democratic Republic of the Congo, and Timor-Leste (United Nations Development Programme, 2009), which produce 67% of the total above ( $\approx 5.2 \times 10^9$  kg), generating almost 600 million International Dollars of wealth per year (United Nations FAO, 2010). Needless to say, a doubling or tripling of below-ground crop tissue due to  $\text{CO}_2$  fertilization would be welcome on both a nutritional and economic basis, especially given that the major grain crops of sub-Saharan Africa are C4 species (sorghum and maize), which show no  $\text{CO}_2$  fertilization response in free-air concentration enrichment (FACE) experiments (Long et al., 2006). Even if some resource limitation is eased by processes inherent to below-ground fertilization, a concomitant increase in water requirement is unavoidable; however, >70% of total water use is already used for irrigated agriculture (Xiong et al., 2010). Locating additional water to support enhanced crop production is unlikely: Ethiopia and Eritrea have among the lowest amount of renewable freshwater resources per capita in Africa (United Nations Statistics Division, 2010). Even though estimates of future  $p\text{CO}_2$  levels imply increased root-crop production, the regions most in need of these benefits may be unable to locate the expanded water resources necessary to truly profit.

## Acknowledgements

We thank E.A. Barth, J.M. Cotton, A.N. Ellenson, A. Greve, W.M. Hagopian, S.A. Harbeson, G.B. Hunsinger, D.P. King, and R.J. Panetta for laboratory assistance. Funding provided by DOE/BES:DE-FG02-09ER16002, by NSF/EAR:0634929, and by the Howard Hughes Foundation.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.agee.2010.11.024.

## References

- Ainsworth, E.A., Long, S.P., 2005. What have we learned from 15 years of free-air  $\text{CO}_2$  enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising  $\text{CO}_2$ . *New Phytol.* 165, 351–372.
- April, R., Keller, D., 1990. Mineralogy of the rhizosphere in forest soils of the Eastern United States: mineralogical studies of the rhizosphere. *Biogeochemistry* 9, 1–18.
- Aranda, I., Pardos, M., Puértolas, J., Jiménez, M.D., Pardos, J.A., 2007. Water-use efficiency in cork oak (*Quercus suber*) is modified by the interaction of water and light availabilities. *Tree Physiol.* 27, 671–677.
- Arens, N.C., Jahren, A.H., Amundson, R., 2000. Can C3 plants faithfully record the carbon isotopic composition of atmospheric carbon dioxide? *Paleobiology* 26, 137–164.
- Baksh, K., Hassan, S., 2005. Use of sewage water for radish cultivation: a case study of Punjab, Pakistan. *J. Agric. Soc. Sci.* 1, 322–326.
- Baldy, P., 1986. Comparison between growth chamber and field-grown *Zea mays* plants for photosynthetic carboxylase activities and other physiological characteristics with respect to leaf position. *J. Exp. Bot.* 37, 309–314.
- Barnes, J.D., Pfirrmann, T., 1992. The influence of  $\text{CO}_2$  and  $\text{O}_3$ , singly and in combination, on gas exchange, growth and nutrient status of radish (*Raphanus sativus* L.). *New Phytol.* 121, 403–412.
- Bhattacharya, N.C., Hileman, D.R., Ghosh, P.P., Musser, R.L., Bhattacharya, S., Biswas, P.K., 1990. Interaction of enriched  $\text{CO}_2$  and water stress on the physiology of and biomass production in sweet potato grown in open-top chambers. *Plant Cell Environ.* 13, 933–940.
- Bowling, D.R., McDowell, N.G., Bond, B.J., Law, B.E., Ehleringer, J.R., 2002.  $^{13}\text{C}$  content of ecosystem respiration is linked to precipitation and vapor pressure deficit. *Oecologia* 131, 113–124.
- Burney, J.A., Davis, S.J., Lobell, D.B., 2010. Greenhouse gas mitigation by agricultural intensification. *Proc. Natl. Acad. Sci. U.S.A.* 107, 12052–12057.
- Cernusak, L.A., Winter, K., Turner, B.L., 2009. Physiological and isotopic ( $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$ ) responses of three tropical tree species to water and nutrient availability. *Plant Cell Environ.* 32, 1441–1455.
- Ceulemans, R., Janssens, I.A., Jach, M.E., 1999. Effects of  $\text{CO}_2$  enrichment on trees and forests: lessons to be learned in view of future ecosystem studies. *Ann. Bot.* 84, 577–590.
- Ceulemans, R., Mousseau, M., 1994. Tansley Review No. 71. Effects of elevated atmospheric  $\text{CO}_2$  on woody plants. *New Phytol.* 127, 425–446.
- Chu, C.C., Coleman, J.S., Mooney, H.A., 1992. Controls of biomass partitioning between roots and shoots: atmospheric  $\text{CO}_2$  enrichment and the acquisition and allocation of carbon and nitrogen in wild radish. *Oecologia* 89, 580–587.
- Conroy, J.P., 1992. Influence of elevated atmospheric  $\text{CO}_2$  concentrations on plant nutrition. *Aust. J. Bot.* 40, 445–456.
- Cotrufo, M.F., Ineson, P., Scott, A., 1998. Elevated  $\text{CO}_2$  reduces the nitrogen concentration of plant tissues. *Global Change Biol.* 4, 43–54.
- Domingos, A.K., Saad, E.B., Wilhelm, H.M., Ramos, L.P., 2008. Optimization of the ethanolation of *Raphanus sativus* (L. Var.) crude oil applying the response surface methodology. *Bioresour. Technol.* 99, 1837–1845.
- Drake, B.G., González-Meler, M.A., Long, S.P., 1997. More efficient plants: a consequence of rising atmospheric  $\text{CO}_2$ ? *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48, 609–639.
- Drever, J.I., 1994. The effect of land plants on weathering rates of silicate minerals. *Geochim. Cosmochim. Acta* 58, 2325–2332.
- Eamus, D., 1991. The interaction of rising  $\text{CO}_2$  and temperatures with water use efficiency. *Plant Cell Environ.* 14, 843–852.
- Edwards, C.E., Haselhorst, M.S.H., McKnite, A.M., Ewers, B.E., Williams, D.G., Weinig, C., 2009. Genotypes of *Brassica rapa* respond differently to plant-induced variation in air  $\text{CO}_2$  concentration in growth chambers with standard and enhanced venting. *Theor. Appl. Genetics* 119, 991–1004.
- Ehleringer, J.R., Cooper, T.A., 1988. Correlations between carbon isotope ratio and microhabitat in desert plants. *Oecologia* 76, 562–566.
- Ehleringer, J.R., Bowling, D.R., Flanagan, L.B., Fessenden, J., Helliiker, B., Martinelli, L.A., Ometto, J.P., 2002. Stable isotopes and carbon cycle processes in forests and grasslands. *Plant Biol.* 4, 181–189.
- Ehleringer, J.R., Field, C.B., Lin, Z.-F., Kuo, C.-Y., 1986. Leaf carbon isotope and mineral composition in subtropical plants along an irradiance cline. *Oecologia* 70, 520–526.
- FAO, 2008. Current World Fertilizer Trends and Outlook to 2011/12. Food and Agriculture Organization of the United Nations.
- Farquhar, G.D., Ehleringer, J.R., Hubick, K.T., 1989. Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40, 503–537.
- Farrar, J.F., Williams, M.L., 1991. The effects of increased atmospheric carbon dioxide and temperature on carbon partitioning, source-sink relations and respiration. *Plant Cell Environ.* 14, 819–830.
- Finzi, A.C., Norby, R.J., Calfapietra, C., Gallet-Budynek, A., Gielen, B., Holmes, W.E., Hoosbeek, M.R., Iversen, C.M., Jackson, R.B., Kubiske, M.E., Ledford, J., Liberloo, M., Oren, R., Polle, A., Pritchard, S., Zak, D.R., Schlesinger, W.H., Ceulemans, R., 2007. Increases in nitrogen uptake rather than nitrogen-use efficiency support higher rates of temperate forest productivity under elevated  $\text{CO}_2$ . *Proc. Natl. Acad. Sci. U.S.A.* 104, 14014–14019.
- Fuhrer, J., 2003. Agroecosystem responses to combinations of elevated  $\text{CO}_2$ , ozone, and global climate change. *Agric. Ecosyst. Environ.* 97, 1–20.
- Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J., Robinson, S., Thomas, S.M., Toulmin, C., 2010. Food security: the challenge of feeding 9 billion people. *Science* 327, 812–818.
- Houghton, J.T., Callander, B.A., Varney, S.K., 1992. Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment. Cambridge University Press, Cambridge.
- Hunt, R., Hand, D.W., Hannah, M.A., Neal, A.M., 1991. Response to  $\text{CO}_2$  enrichment in 27 herbaceous species. *Funct. Ecol.* 5, 410–421.
- Hunt, R., Hand, D.W., Hannah, M.A., Neal, A.M., 1993. Further responses to  $\text{CO}_2$  enrichment in British herbaceous species. *Funct. Ecol.* 7, 661–668.
- Idso, S.B., Idso, K.E., 2001. Effects of atmospheric  $\text{CO}_2$  enrichment on plant constituents related to animal and human health. *Environ. Exp. Bot.* 45, 179–199.
- Idso, S.B., Kimball, B.A., 1989. Growth response of carrot and radish to atmospheric  $\text{CO}_2$  enrichment. *Environ. Exp. Bot.* 29, 135–139.
- Idso, S.B., Kimball, B.A., Mauney, J.R., 1988. Effects of atmospheric  $\text{CO}_2$  enrichment on root:shoot ratios of carrot, radish, cotton and soybean. *Agric. Ecosyst. Environ.* 21, 293–299.
- Jablonski, L.M., 1997. Responses of vegetative and reproductive traits to elevated  $\text{CO}_2$  and nitrogen in *Raphanus* varieties. *Can. J. Bot.* 75, 533–545.
- Jackson, R.B., Sala, O.E., Field, C.B., Mooney, H.A., 1994.  $\text{CO}_2$  alters water use, carbon gain, and yield for the dominant species in a natural grassland. *Oecologia* 98, 257–262.
- Jahren, A.H., Arens, N.C., Harbeson, S.A., 2008. Prediction of atmospheric  $\delta^{13}\text{C}$  using fossil plant tissues. *Rev. Geophys.* 46, 1–12.
- Javald, A., 2009. Arbuscular mycorrhizal mediated nutrition in plants. *J. Plant Nutr.* 32, 1595–1618.
- Kates, R.W., Dasgupta, P., 2007. African poverty: a grand challenge for sustainability science. *Proc. Natl. Acad. Sci. U.S.A.* 104, 16747–16750.



- Kimball, B.A., Mauney, J.R., Nakamura, H., Idso, S.B., 1993. Effects of increasing atmospheric CO<sub>2</sub> on vegetation. *Vegetatio* 104/105, 65–75.
- Knecht, G.N., 1975. Response of radish to high CO<sub>2</sub>. *HortScience* 10, 274–275.
- Kostka-Rick, R., Manning, W.J., 1993. Radish (*Raphanus sativus* L.): a model for studying responses to air pollutants and other environmental stresses. *Environ. Pollut.* 82, 107–138.
- Kranabetter, J.M., Simard, S.W., Guy, R.D., Coates, K.D., 2010. Species patterns in foliar nitrogen concentration, nitrogen content and <sup>13</sup>C abundance for understory saplings across light gradients. *Plant Soil* 327, 389–401.
- Langley, J.A., Megonigal, J.P., 2010. Ecosystem response to elevated CO<sub>2</sub> levels limited by nitrogen-induced plant species shift. *Nature* 466, 96–99.
- Lawlor, D.W., Mitchell, R.A.C., 1991. The effects of increasing CO<sub>2</sub> on crop photosynthesis and productivity: a review of field studies. *Plant Cell Environ.* 14, 807–818.
- Leadley, P.W., Drake, B.G., 1993. Open top chambers for exposing plant canopies to elevated CO<sub>2</sub> concentration and for measuring net gas exchange. *Vegetatio* 104/105, 3–15.
- Li, P., Ainsworth, E.A., Leakey, A.D.B., Ulanov, A., Lozovaya, V., Ort, D.R., Bohnert, H.J., 2008. Arabidopsis transcript and metabolite profiles: ecotype-specific responses to open-air elevated [CO<sub>2</sub>]. *Plant Cell Environ.* 31, 1673–1687.
- Lin, W., Zhang, F., Bai, K., 2000. Responses of plant rhizosphere to atmospheric CO<sub>2</sub> enrichment. *Chin. Sci. Bull.* 45, 97–101.
- Lloyd, J., Farquhar, G.D., 1996. The CO<sub>2</sub> dependence of photosynthesis, plant growth responses to elevated atmospheric CO<sub>2</sub> concentrations and their interaction with soil nutrient status. I. General principles and forest ecosystems. *Funct. Ecol.* 10, 4–32.
- Lobell, D.B., Field, C.B., 2008. Estimation of the carbon dioxide (CO<sub>2</sub>) fertilization effect using growth rate anomalies of CO<sub>2</sub> and crop yields since 1961. *Global Change Biol.* 14, 39–45.
- Long, S.P., Ainsworth, E.A., Leakey, A.D.B., Nösberger, J., Ort, D.R., 2006. Food for thought: lower-than-expected crop yield stimulation with rising CO<sub>2</sub> concentrations. *Science* 312, 1918–1921.
- Lord, D., Morissette, S., Allaire, J., 1993. Influence of light intensity, nocturnal air temperature and carbon dioxide levels on greenhouse black spruce seedlings (*Picea mariana*). *Can. J. For. Res.* 23, 101–110.
- Lucier, G., Jerardo, A., 2004. Electronic Outlook Report from the Economic Research Service: Vegetables and Melons Outlook. USDA.
- Ludlow, M.M., Jarvis, P.G., 1971. Methods for measuring photorespiration in leaves. In: Šestak, Z., Čatský, J., Jarvis, P.G. (Eds.), *Plant Photosynthetic Production: Manual of Methods*. Dr W. Junk N.V. Publishers, The Hague, pp. 294–315.
- Luxmoore, R.J., Oneill, E.G., Ells, J.M., Rogers, H.H., 1986. Nutrient uptake and growth responses of Virginia pine to elevated atmospheric carbon dioxide. *J. Environ. Qual.* 15, 244–251.
- Lyon, T.L., Buckman, H.O., Brady, N.C., 1952. *The Nature and Properties of Soils*. The Macmillan Company, New York.
- Medlyn, B.E., Barton, C.V.M., Broadmeadow, M.S.J., Ceulemans, R., Angelis, P.D., Forstreuter, M., Freeman, M., Jackson, S.B., Kellomäki, S., Laitat, E., Rey, A., Robertz, P., Sigurdsson, B.D., Strassmeyer, J., Wang, K., Curtis, P.S., Jarvis, P.G., 2001. Stomatal conductance of forest species after long-term exposure to elevated CO<sub>2</sub> concentration: a synthesis. *New Phytol.* 149, 247–264.
- Melillo, J.M., McGuire, A.D., Kicklighter, D.W., Moore III, B., Vorosmarty, C.J., Schloss, A.L., 1993. Global climate change and terrestrial net primary production. *Nature* 363, 234–240.
- Miglietta, F., Bindi, M., Vaccai, F.P., Schapendonk, A.H.C.M., Wolf, J., Butterfield, R.E., 2000. Crop ecosystem responses to climatic change: root and tuberous crops. In: Reddy, K.R., Hodges, H.F. (Eds.), *Climate Change and Global Crop Productivity*. CABI Publishing, Wallingford, UK, pp. 189–212.
- Morison, J.I.L., Gifford, R.M., 1984. Plant growth and water use with limited water supply in high CO<sub>2</sub> concentrations. II. Plant dry weight, partitioning and water use efficiency. *Aust. J. Plant Physiol.* 11, 375–384.
- Nakicenovic, N., Swart, R., 2000. *Emission Scenarios: Summary for Policymakers*. Cambridge University Press, Cambridge.
- NOAA/ESRL, Dr. Pieter Tans, Trends in Atmospheric Carbon Dioxide. [www.esrl.noaa.gov/gmd/ccgg/trends/](http://www.esrl.noaa.gov/gmd/ccgg/trends/) (accessed: July 7, 2010).
- Norby, R.J., Leford, J., Reilly, C.D., Miller, N.E., O'Neill, E.G., 2004. Fine-root production dominates response of a deciduous forest to atmospheric CO<sub>2</sub> enrichment. *Proc. Natl. Acad. Sci. U.S.A.* 101, 9689–9693.
- Nowak, R.S., Ellsworth, D.S., Smith, S.D., 2004. Functional responses of plants to elevated atmospheric CO<sub>2</sub>—do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytol.* 162, 253–280.
- Otten, J.J., Hellwig, J.P., Meyers, L.D., 2006. *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. The National Academies Press, Washington, DC.
- Overdieck, D., Reid, C., Strain, B.R., 1988. The effects of preindustrial and future CO<sub>2</sub> concentrations on growth, dry matter production and the C/N relationship in plants at low nutrient supply: *Vigna unguiculata* (cowpea), *Abelmoschus esculentus* (okra) and *Raphanus sativus* (radish). *Angew. Bot.* 62, 119–134.
- Overdieck, D., Reining, E., 1986. Effect of atmospheric CO<sub>2</sub> enrichment on perennial ryegrass (*Lolium perenne* L.) and white clover (*Trifolium repens* L.) competing in managed model-ecosystems. II. Nutrient uptake. *Acta Oecol. Oecol. Plant.* 7, 367–378.
- Park, R., Epstein, S., 1960. Carbon isotope fractionation during photosynthesis. *Geochim. Cosmochim. Acta* 21, 110–126.
- Poorter, H., 1993. Interspecific variation in the growth response of plants to an elevated ambient CO<sub>2</sub> concentration. *Vegetatio* 104/105, 77–97.
- Poorter, H., Garnier, E., 1996. Plant growth analysis: an evaluation of experimental design and computational methods. *J. Exp. Bot.* 47, 1343–1351.
- Poorter, H., Navas, M.-L., 2003. Plant growth and competition at elevated CO<sub>2</sub>: on winners, losers and functional groups. *New Phytol.* 157, 175–198.
- Riechers, G.H., Strain, B.R., 1988. Growth response of blue grama (*Bouteloua gracilis*) in response to atmospheric CO<sub>2</sub> enrichment. *Can. J. Bot.* 66, 1570–1573.
- Rogers, H.H., Peterson, C.M., McCrimmon, J.N., Cure, J.D., 1992. Resonance of plant roots to elevated atmospheric carbon dioxide. *Plant Cell Environ.* 15, 749–752.
- Rogers, H.H., Prior, S.A., Runion, G.B., Mitchell, R.J., 1996. Root to shoot ratio of crops as influenced by CO<sub>2</sub>. *Plant Soil* 187, 229–248.
- Rogers, H.H., Runion, G.B., Krupa, S.V., 1994. Plant responses to atmospheric CO<sub>2</sub> enrichment with emphasis on roots and the rhizosphere. *Environ. Pollut.* 83, 155–189.
- Schenk, U., Jäger, H.-J., Weigel, H.-J., 1996. Nitrogen supply determines responses of yield and biomass partitioning of perennial ryegrass to elevated atmospheric carbon dioxide concentrations. *J. Plant Nutr.* 19, 1423–1440.
- Schmidhuber, J., Tubiello, F.N., 2007. Global food security under climate change. *Proc. Natl. Acad. Sci. U.S.A.* 104, 19703–19708.
- Schwarz, N., Strain, B.R., 1990. Carbon—a plant nutrient, deficiency and sufficiency. *J. Plant Nutr.* 13, 1073–1078.
- Sionit, N., 1983. Response of soybean to two levels of mineral nutrition in CO<sub>2</sub>-enriched atmosphere. *Crop Sci.* 23, 329–333.
- Sionit, N., Hellmers, H., Strain, B.R., 1982. Interaction of atmospheric CO<sub>2</sub> enrichment and irradiance on plant growth. *Agron. J.* 74, 721–725.
- Snow, A.A., Campbell, L.G., 2005. Can feral radishes become weeds? In: Gressel, J. (Ed.), *Crop Fertility and Volunteerism*. CRC Press, Taylor and Francis Group, LLC, Boca Raton, FL, pp. 193–208.
- Soil Survey Staff, Soil Conservation Service, 1999. *Keys to Soil Taxonomy*. Pocahontas Press, Inc., VA.
- Stuhlfauth, T., Klug, K., Fock, H.P., 1987. The production of secondary metabolites by *Digitalis lanata* during CO<sub>2</sub> enrichment and water stress. *Phytochemistry* 26, 2735–2739.
- Taub, D.R., Miller, B., Allen, H., 2008. Effects of elevated CO<sub>2</sub> on the protein concentration of food crops: a meta-analysis. *Global Change Biol.* 14, 565–575.
- Taub, D.R., Wang, X., 2008. Why are nitrogen concentrations in plant tissues lower under elevated CO<sub>2</sub>? A critical examination of the hypotheses. *J. Integr. Plant Biol.* 50, 1365–1374.
- Tognoni, F., Halevy, A.H., Wittwer, S.H., 1966. Growth of bean and tomato plants as affected by root absorbed growth substances and atmospheric carbon dioxide. *Planta* 72, 43–52.
- Tubiello, F.N., Soussana, J.-F., Howden, S.M., 2007. Crop and pasture response to climate change. *Proc. Natl. Acad. Sci. U.S.A.* 104, 19686–19690.
- United Nations Development Programme, 2009. *Human Development Report 2009. Overcoming Barriers: Human Mobility and Development*. Palgrave Macmillan, New York.
- United Nations FAO, Food and Agriculture Organization of the United Nations. <http://faostat.fao.org/site/339/default.aspx> (accessed: June 22, 2010).
- United Nations MDGI, United Nations Millennium Development Goals Indicators. <http://mdgs.un.org/unsd/mdg/SeriesDetail.aspx?sr=749&crd=6> (accessed: June 22, 2010).
- United Nations Statistics Division, Environmental Indicators, Water. <http://unstats.un.org/unsd/ENVIRONMENT/waterresources.htm> (accessed: June 22, 2010).
- van Hees, P.A.W., Rosling, A., Lundström, U.S., Finlay, R.D., 2006. The biogeochemical impact of ectomycorrhizal conifers on major soil elements (Al, Fe, K and Si). *Geoderma* 136, 364–377.
- Vessey, J.K., Henry, L.T., Raper, C.D., 1990. Nitrogen nutrition and temporal effects of enhanced carbon dioxide on soybean growth. *Crop Sci.* 30, 287–294.
- Wand, S.J.E., Midgley, G.F., Jones, M.H., Curtis, P.S., 1999. Responses of wild C4 and C3 grass (Poaceae) species to elevated atmospheric CO<sub>2</sub> concentration: a meta-analytic test of current theories and perceptions. *Global Change Biol.* 5, 723–741.
- Wang, B., Qiu, Y.L., 2006. Phylogenetic distribution and evolution of mycorrhizas in land plants. *Mycorrhiza* 16, 299–363.
- Wheeler, R.M., Mackowiak, C.L., Sager, J.C., Knott, W.M., 1994. Growth of soybean and potato at high CO<sub>2</sub> partial pressures. *Adv. Space Res.* 14, 251–255.
- Wheeler, R.M., Tibbitts, T.W., 1989. Utilization of potatoes for life support systems in space. IV. Effect of CO<sub>2</sub> enrichment. *Am. J. Potato Res.* 66, 25–34.
- Wheeler, T.R., Morison, J.I.L., Ellis, R.H., Hadley, P., 1994. The effects of CO<sub>2</sub>, temperature and their interaction on the growth and yield of carrot (*Daucus carota* L.). *Plant Cell Environ.* 17, 1275–1284.
- Wong, S.-C., 1993. Interaction between elevated atmospheric concentration of CO<sub>2</sub> and humidity on plant growth: comparison between cotton and radish. *Vegetatio* 104/105, 211–221.
- Wong, S.C., 1979. Elevated atmospheric partial pressure of CO<sub>2</sub> and plant growth. I. Interactions of nitrogen nutrition and photosynthetic capacity in C<sub>3</sub> and C<sub>4</sub> plants. *Oecologia* 44, 68–74.
- Xiong, W., Holman, I., Lin, E., Conway, D., Jiang, J., Xu, Y., Li, Y., 2010. Climate change, water availability and future cereal production in China. *Agric. Ecosyst. Environ.* 135, 58–69.
- Yin, C.Y., Pang, X.Y., Chen, K., 2009. The effects of water, nutrient availability and their interaction on the growth, morphology and physiology of two poplar species. *Environ. Exp. Bot.* 67, 196–203.

- Yin, C.Y., Wang, X., Duan, B.L., Luo, J.X., Li, C.Y., 2005. Early growth, dry matter allocation and water use efficiency of two sympatric *Populus* species as affected by water stress. *Environ. Exp. Bot.* 53, 315–322.
- Zachos, J.C., Dickens, G.R., Zeebe, R.E., 2008. An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* 451, 279–283.
- Zangerl, A.R., Bazzaz, F.A., 1984. The response of plants to elevated CO<sub>2</sub>. II. Competitive interactions among annual plants under varying light and nutrients. *Oecologia* 62, 412–417.
- Zeng, R.S., Mallik, A.U., Setliff, E., 2003. Growth stimulation of ectomycorrhizal fungi by root exudates of Brassicaceae plants: role of degraded compounds of indole glucosinolates. *J. Chem. Ecol.* 29, 1337–1355.
- Zhang, F., Shen, J., Zhang, J., Zuo, Y., Li, L., Chen, X., 2010. Rhizosphere processes and management for improving nutrient use efficiency and crop productivity: implications for China. *Adv. Agron.* 107, 1–32.
- Zimmerman, J.K., Ehleringer, J.R., 1990. Carbon isotope ratios are correlated with irradiance levels in the Panamanian orchid *Catasetum viridiflavum*. *Oecologia* 83, 247–249.



**Table S1**Elevated  $p\text{CO}_2$  experiments performed.<sup>a</sup>

$p\text{CO}_2$ (ppmv) <sup>b</sup>	Temperature (°C) <sup>b</sup>	Humidity (%) <sup>b</sup>
348 ± 17	23 ± 1	59 ± 6
348 ± 17	29 ± 2	40 ± 6
388 ± 26	23 ± 1	36 ± 8
388 ± 26	29 ± 1	27 ± 4
413 ± 39	23.9 ± 1.5	23 ± 3
413 ± 39	30.0 ± 1.8	20 ± 3
426	22.5	– <sup>c</sup>
426	29.7	–
760 ± 30	23 ± 1	32 ± 8
760 ± 30	29 ± 1	27 ± 5
1090 ± 30	23 ± 1	53 ± 5
1090 ± 30	29 ± 2	38 ± 7
1425 ± 59	22.8 ± 1.1	21 ± 2
1425 ± 59	29.7 ± 1.8	19 ± 3
1791	25.5	–
1791	30.1	–

<sup>a</sup>Eight plants were grown for each set of conditions, with the exception of  $p\text{CO}_2 = 413$  and 1425 ppmv for which 9 plants were grown in each experiment.

<sup>b</sup>All values are reported as the mean ± 1σ, as measured throughout the four-week experiment.

<sup>c</sup>Not measured.

**Table S2**  
Percent biomass increase data.

Species	Common name	Biomass part <sup>a</sup>	Control pCO <sub>2</sub> (ppmv)	Elevated pCO <sub>2</sub> (ppmv)	% biomass increase	Reference <sup>c</sup>
<i>Beta vulgaris</i>	sugar beet	AG	332	725	49	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	AG	332	725	153	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	AG	332	725	28	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	-11	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	5	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	32	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	20	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	5	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	4	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	AG	375	550	21	(Manderscheid et al., 2010)
<i>Ipomoea batatas</i>	sweet potato	AG	364	438	6	(Bhattacharya et al., 1990)
<i>Ipomoea batatas</i>	sweet potato	AG	364	666	9	(Bhattacharya et al., 1990)
<i>Ipomoea batatas</i>	sweet potato	AG	354	431	22	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	AG	354	506	23	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	AG	354	659	12	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	AG	354	438	-15	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	AG	354	514	-2	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	AG	354	665	19	(Biswas et al., 1996)
<i>Raphanus sativus</i>	radish	AG	386	761	7	(Barnes and Pfirrmann, 1992)
<i>Raphanus sativus</i>	radish	AG	330	600	25	(Chu et al., 1992)
<i>Raphanus sativus</i>	radish	AG	350	700	8	(Wong, 1993)
<i>Raphanus sativus</i>	radish	AG	350	700	52	(Wong, 1993)
<i>Raphanus sativus</i>	radish	AG	350	650	22	(Overdieck et al., 1988)
<i>Raphanus sativus</i>	radish	AG	400	1200	66	(Knecht, 1975)
<i>Solanum tuberosum</i>	potato	AG	380	550	-8	(Högy and Fangmeier, 2009)
<i>Solanum tuberosum</i>	potato	AG	ambient <sup>b</sup>	680	-43	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	-26	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	-10	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	-12	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	-15	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	26	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	-4	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	-2	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	ambient	680	28	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	AG	398	543	23	(Donnelly et al., 2001)
<i>Solanum tuberosum</i>	potato	AG	398	694	28	(Donnelly et al., 2001)
<i>Solanum tuberosum</i>	potato	AG	369	543	-16	(Conn and Cochran, 2006)
<i>Solanum tuberosum</i>	potato	AG	369	707	0	(Conn and Cochran, 2006)
<i>Beta vulgaris</i>	sugar beet	BG	332	775	29	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	BG	332	775	178	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	BG	332	775	87	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	BG	375	550	14	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	BG	375	550	10	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	BG	375	550	17	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	BG	375	550	8	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	BG	375	550	36	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	BG	375	550	18	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	BG	378	668	26	(Demmers-Derks et al., 1998)
<i>Daucus carota</i>	carrot	BG	348	551	31	(Wheeler et al., 1994)
<i>Daucus carota</i>	carrot	BG	348	551	80	(Wheeler et al., 1994)
<i>Ipomoea batatas</i>	sweet potato	BG	364	438	63	(Bhattacharya et al., 1990)
<i>Ipomoea batatas</i>	sweet potato	BG	364	666	40	(Bhattacharya et al., 1990)
<i>Ipomoea batatas</i>	sweet potato	BG	361	438	26	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	BG	361	514	43	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	BG	361	665	69	(Biswas et al., 1996)
<i>Raphanus sativus</i>	radish	BG	350	650	70	(Jablonski, 1997)
<i>Raphanus sativus</i>	radish	BG	330	600	40	(Chu et al., 1992)
<i>Raphanus sativus</i>	radish	BG	386	761	43	(Barnes and Pfirrmann, 1992)
<i>Raphanus sativus</i>	radish	BG	350	700	289	(Wong, 1993)
<i>Raphanus sativus</i>	radish	BG	350	700	113	(Wong, 1993)
<i>Raphanus sativus</i>	radish	BG	350	650	78	(Overdieck et al., 1988)
<i>Solanum tuberosum</i>	potato	BG	360	460	-5	(Miglietta et al., 1998)
<i>Solanum tuberosum</i>	potato	BG	360	560	9	(Miglietta et al., 1998)
<i>Solanum tuberosum</i>	potato	BG	360	660	4	(Miglietta et al., 1998)
<i>Solanum tuberosum</i>	potato	BG	380	550	6	(Högy and Fangmeier, 2009)

<i>Solanum tuberosum</i>	potato	BG	370	715	29	(Heagle et al., 2003)
<i>Solanum tuberosum</i>	potato	BG	370	540	20	(Heagle et al., 2003)
<i>Solanum tuberosum</i>	potato	BG	370	715	22	(Heagle et al., 2003)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	3	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	16	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	24	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	9	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	12	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	40	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	54	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	-7	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	11	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	ambient	680	41	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	BG	398	543	26	(Donnelly et al., 2001)
<i>Solanum tuberosum</i>	potato	BG	398	694	41	(Donnelly et al., 2001)
<i>Solanum tuberosum</i>	potato	BG	369	543	20	(Conn and Cochran, 2006)
<i>Solanum tuberosum</i>	potato	BG	369	707	36	(Conn and Cochran, 2006)
<i>Beta vulgaris</i>	sugar beet	TOT	350	675	94	(Sionit et al., 1982)
<i>Beta vulgaris</i>	sugar beet	TOT	350	675	114	(Sionit et al., 1982)
<i>Beta vulgaris</i>	sugar beet	TOT	366	703	55	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	TOT	332	725	41	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	TOT	332	725	141	(Ziska et al., 1995)
<i>Beta vulgaris</i>	sugar beet	TOT	375	550	6	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	TOT	375	550	8	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	TOT	375	550	18	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	TOT	375	550	7	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	TOT	375	550	29	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	TOT	375	550	19	(Manderscheid et al., 2010)
<i>Beta vulgaris</i>	sugar beet	TOT	372	592	16	(Demmers-Derks et al., 1998)
<i>Beta vulgaris</i>	sugar beet	TOT	377	709	27	(Demmers-Derks et al., 1998)
<i>Beta vulgaris</i>	sugar beet	TOT	384	704	25	(Demmers-Derks et al., 1998)
<i>Daucus carota</i>	carrot	TOT	340	640	176	(Idso and Kimball, 1989)
<i>Daucus carota</i>	carrot	TOT	348	551	16	(Wheeler et al., 1994)
<i>Daucus carota</i>	carrot	TOT	348	551	25	(Wheeler et al., 1994)
<i>Ipomoea batatas</i>	sweet potato	TOT	364	666	46	(Bhattacharya et al., 1990)
<i>Ipomoea batatas</i>	sweet potato	TOT	364	666	31	(Bhattacharya et al., 1990)
<i>Ipomoea batatas</i>	sweet potato	TOT	361	438	2	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	TOT	361	514	17	(Biswas et al., 1996)
<i>Ipomoea batatas</i>	sweet potato	TOT	361	665	40	(Biswas et al., 1996)
<i>Raphanus sativus</i>	radish	TOT	340	680	72	(Morison and Gifford, 1984)
<i>Raphanus sativus</i>	radish	TOT	350	650	50	(Jablonski, 1997)
<i>Raphanus sativus</i>	radish	TOT	330	600	30	(Chu et al., 1992)
<i>Raphanus sativus</i>	radish	TOT	340	640	52	(Idso and Kimball, 1989)
<i>Raphanus sativus</i>	radish	TOT	350	675	63	(Sionit et al., 1982)
<i>Raphanus sativus</i>	radish	TOT	350	675	33	(Sionit et al., 1982)
<i>Raphanus sativus</i>	radish	TOT	386	761	36	(Barnes and Pfirrmann, 1992)
<i>Raphanus sativus</i>	radish	TOT	350	700	73	(Wong, 1993)
<i>Raphanus sativus</i>	radish	TOT	350	700	67	(Wong, 1993)
<i>Raphanus sativus</i>	radish	TOT	350	650	51	(Overdieck et al., 1988)
<i>Solanum tuberosum</i>	potato	TOT	380	550	5	(Högy and Fangmeier, 2009)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	-3	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	3	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	18	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	7	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	9	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	37	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	39	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	12	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	ambient	680	28	(Craigon et al., 2002)
<i>Solanum tuberosum</i>	potato	TOT	398	543	25	(Donnelly et al., 2001)
<i>Solanum tuberosum</i>	potato	TOT	398	694	38	(Donnelly et al., 2001)

<sup>a</sup>AG = above ground, BG = below ground, TOT = Total

<sup>b</sup>ambient =  $p\text{CO}_2$  level for control chamber not given.

<sup>c</sup>References are as follows:

Barnes, J.D., Pfirrmann, T., 1992. The influence of  $\text{CO}_2$  and  $\text{O}_3$ , singly and in combination, on gas exchange, growth and nutrient status of radish (*Raphanus sativus* L.). New Phytol. 121, 403-412.

- Bhattacharya, N.C., Hileman, D.R., Ghosh, P.P., Musser, R.L., Bhattacharya, S., Biswas, P.K., 1990. Interaction of enriched CO<sub>2</sub> and water stress on the physiology of and biomass production in sweet potato grown in open-top chambers. *Plant, Cell Environ.* 13, 933-940.
- Biswas, P.K., Hileman, D.R., Ghosh, P.P., Bhattacharya, N.C., McCrimmon, J.N., 1996. Growth and yield responses of field-grown sweetpotato to elevated carbon dioxide. *Crop. Sci.* 36, 1234-1239.
- Chu, C.C., Coleman, J.S., Mooney, H.A., 1992. Controls of biomass partitioning between roots and shoots: Atmospheric CO<sub>2</sub> enrichment and the acquisition and allocation of carbon and nitrogen in wild radish. *Oecol.* 89, 580-587.
- Conn, J.S., Cochran, V.L., 2006. Response of potato (*Solanum tuberosum* L.) to elevated atmospheric CO<sub>2</sub> in the North American subarctic. *Agric., Ecosyst. Environ.* 112, 49-57.
- Craigon, J., Fangmeier, A., Jones, M., Donnelly, A., Bindi, M., De Temmerman, L., Persson, K., Ojanpera, K., 2002. Growth and marketable-yield responses of potato to increased CO<sub>2</sub> and ozone. *Eur. J. Agron.* 17, 273-289.
- Demmers-Derks, H., Mitchell, R.A.C., Mitchell, V.J., Lawlor, D.W., 1998. Response of sugar beet (*Beta vulgaris* L.) yield and biochemical composition to elevated CO<sub>2</sub> and temperature at two nitrogen applications. *Plant Cell. Environ.* 21, 829-836.
- Donnelly, A., Craigon, J., Black, C.R., Colls, J.J., Landon, G., 2001. Elevated CO<sub>2</sub> increases biomass and tuber yield in potato even at high ozone concentrations. *New Phytol.* 149, 265-274.
- Heagle, A.S., Miller, J.E., Pursley, W.A., 2003. Growth and yield responses of potato to mixtures of carbon dioxide and ozone. *J. Environ. Qual.* 32, 1603-1610.
- Högy, P., Fangmeier, A., 2009. Atmospheric CO<sub>2</sub> enrichment affects potatoes: 1. Aboveground biomass production and tuber yield. *Eur. J. Agron.* 30, 78-84.
- Idso, S.B., Kimball, B.A., 1989. Growth response of carrot and radish to atmospheric CO<sub>2</sub> enrichment. *Environ. Exp. Bot.* 29, 135-139.
- Jablonski, L.M., 1997. Responses of vegetative and reproductive traits to elevated CO<sub>2</sub> and nitrogen in *Raphanus* varieties. *Can. J. Bot.* 75, 533-545.
- Knecht, G.N., 1975. Response of radish to high CO<sub>2</sub>. *HortSci.* 10, 274-275.
- Manderscheid, R., Pacholski, A., Weigel, H.-J., 2010. Effect of free air carbon dioxide enrichment combined with two nitrogen levels on growth, yield and yield quality of sugar beet: Evidence for a sink limitation of beet growth under elevated CO<sub>2</sub>. *Eur. J. Agron.* 32, 228-239.
- Miglietta, F., Magliulo, V., Bindi, M., Cerio, L., Vaccari, F.P., Loduca, V., Peressotti, A., 1998. Free air CO<sub>2</sub> enrichment of potato (*Solanum tuberosum* L.): development, growth and yield. *Global Change Biol.* 4, 163-172.
- Morison, J.I.L., Gifford, R.M., 1984. Plant growth and water use with limited water supply in high CO<sub>2</sub> concentrations. II. Plant dry weight, partitioning and water use efficiency. *Aust. J. Plant Physiol.* 11, 375-384.
- Overdieck, D., Reid, C., Strain, B.R., 1988. The effects of preindustrial and future CO<sub>2</sub> concentrations on growth, dry matter production and the C/N relationship in plants at low nutrient supply: *Vigna unguiculata* (cowpea), *Abelmoschus esculentus* (okra) and *Raphanus sativus* (radish). *Angew. Bot.* 62, 119-134.



- Sionit, N., Hellmers, H., Strain, B.R., 1982. Interaction of atmospheric CO<sub>2</sub> enrichment and irradiance on plant growth. *Agron. J.* 74, 721-725.
- Wheeler, T.R., Morison, J.I.L., Ellis, R.H., Hadley, P., 1994. The effects of CO<sub>2</sub>, temperature and their interaction on the growth and yield of carrot (*Daucus carota* L.). *Plant Cell. Environ.* 17, 1275-1284.
- Wong, S.-C., 1993. Interaction between elevated atmospheric concentration of CO<sub>2</sub> and humidity on plant growth: comparison between cotton and radish. *Vegetatio* 104/105, 211-221.
- Ziska, L.H., Sicher, R.C., Kremer, D.F., 1995. Reversibility of photosynthetic acclimation of swiss chard and sugarbeet grown at elevated concentrations of CO<sub>2</sub>. *Physiol. Plant.* 95, 355-364.

**Table S3**  
Biomass data.

Plant #	$p\text{CO}_2$ (ppmv)	Above ground [ln(mg)] <sup>a</sup>	Below ground [ln(mg)] <sup>a</sup>
1	348	5.39	3.91
2	348	5.08	2.30
3	348	4.94	2.30
4	348	5.39	3.91
5	348	5.63	4.94
6	348	5.14	4.09
7	348	4.70	3.91
8	348	4.61	2.30
9	348	5.25	3.69
10	348	5.19	3.69
11	348	5.19	3.00
12	348	5.44	3.00
13	348	5.19	4.50
14	348	4.87	4.94
15	348	5.30	4.50
16	348	5.25	4.50
<i>Average</i>	348	$5.19 \pm 0.27$	$4.04 \pm 0.90$
17	388	5.52	3.91
18	388	5.60	5.19
19	388	5.25	5.25
20	388	4.79	2.30
21	388	5.39	5.30
22	388	5.56	5.30
23	388	5.70	4.38
24	388	5.67	5.14
25	388	5.35	3.91
26	388	5.60	3.91
27	388	5.56	2.30
28	388	4.79	2.30
29	388	5.01	3.91
30	388	3.91	4.09
31	388	5.48	2.30
32	388	5.44	– <sup>b</sup>
<i>Average</i>	388	$5.37 \pm 0.47$	$4.48 \pm 1.18$
33	413	5.46	5.59
34	413	5.08	4.32
35	413	6.07	5.68
36	413	5.14	5.13
37	413	5.46	5.41
38	413	5.48	5.07
39	413	5.79	4.75
40	413	5.68	5.39
41	413	5.03	2.77
42	413	5.38	3.18
43	413	5.23	4.53
44	413	5.67	4.91
45	413	5.73	5.56
46	413	5.59	5.24
47	413	5.98	4.97
48	413	5.64	3.67
49	413	5.43	3.75
50	413	5.84	3.46
<i>Average</i>	413	$5.58 \pm 0.30$	$4.93 \pm 0.90$
51	426	5.59	5.63
52	426	5.54	5.93
53	426	5.61	5.79
54	426	5.66	6.10
55	426	5.48	6.17
56	426	5.88	4.42
57	426	5.16	5.27
58	426	5.64	4.33
59	426	5.84	5.63
60	426	5.59	5.61
61	426	5.61	5.38
62	426	5.32	5.91
63	426	5.65	5.40
64	426	5.13	5.07
65	426	5.72	4.88
66	426	5.73	5.07
<i>Average</i>	426	$5.59 \pm 0.21$	$5.54 \pm 0.55$

Plant #	$p\text{CO}_2$ (ppmv)	Above ground [ln(mg)]	Below ground [ln(mg)]
67	760	5.56	5.80
68	760	5.77	6.02
69	760	5.60	6.11
70	760	5.83	6.02
71	760	6.02	6.13
72	760	5.67	5.74
73	760	6.09	6.13
74	760	5.60	5.74
75	760	5.60	5.60
76	760	5.52	4.50
77	760	5.86	5.08
78	760	5.80	5.77
79	760	5.35	5.52
80	760	5.56	4.38
81	760	6.09	5.35
82	760	5.60	5.74
<i>Average</i>	760	$5.74 \pm 0.21$	$5.71 \pm 0.54$
83	1090	5.67	5.56
84	1090	5.60	5.63
85	1090	5.60	6.29
86	1090	5.63	4.94
87	1090	6.04	5.08
88	1090	5.44	5.52
89	1090	5.63	5.83
90	1090	5.35	6.23
91	1090	4.94	4.94
92	1090	6.02	5.39
93	1090	6.06	5.35
94	1090	5.63	4.25
95	1090	5.67	5.48
96	1090	5.60	3.91
97	1090	5.83	4.38
98	1090	5.39	4.61
<i>Average</i>	1090	$5.67 \pm 0.28$	$5.41 \pm 0.68$
99	1425	6.00	6.48
100	1425	6.04	5.66
101	1425	6.13	6.06
102	1425	5.73	6.04
103	1425	5.94	5.76
104	1425	5.74	5.23
105	1425	5.81	5.29
106	1425	5.54	5.17
107	1425	5.60	5.09
108	1425	5.97	5.16
109	1425	5.91	5.39
110	1425	5.51	5.50
111	1425	5.38	5.59
112	1425	5.78	5.68
113	1425	5.99	5.31
114	1425	5.72	3.59
115	1425	5.86	5.24
116	1425	5.55	3.02
<i>Average</i>	1425	$5.81 \pm 0.21$	$5.52 \pm 0.82$
117	1791	5.87	6.20
118	1791	5.99	–
119	1791	5.90	6.42
120	1791	5.42	6.42
121	1791	5.45	6.07
122	1791	5.65	6.30
123	1791	5.65	6.33
124	1791	5.18	5.98
125	1791	5.39	5.50
126	1791	5.72	5.58
127	1791	5.84	6.02
128	1791	5.86	6.36
129	1791	5.90	5.77
130	1791	5.89	4.18
131	1791	5.74	6.30
132	1791	6.10	5.20
<i>Average</i>	1791	$5.75 \pm 0.25$	$5.87 \pm 0.60$

<sup>a</sup>All reported uncertainties are  $1\sigma$ .

<sup>b</sup>No data.