

Variation in oxygen isotope fractionation during cellulose synthesis: intramolecular and biosynthetic effects

LEONEL STERNBERG¹, MARIA CAMILA PINZON¹, WILLIAM T. ANDERSON² & A. HOPE JAHREN³

¹Department of Biology, University of Miami, Coral Gables, Fl. 33124, USA, ²Earth Sciences Department/Southeast Environmental Research Center, Florida International University, 11200 S.W. 8 St. Miami, FL 33199, USA and ³Earth and Planetary Sciences, John Hopkins University, Baltimore, MD, 21218, USA

ABSTRACT

The oxygen isotopic composition of plant cellulose is commonly used for the interpretations of climate, ecophysiology and dendrochronology in both modern and palaeoenvironments. Further applications of this analytical tool depends on our in-depth knowledge of the isotopic fractionations associated with the biochemical pathways leading to cellulose. Here, we test two important assumptions regarding isotopic effects resulting from the location of oxygen in the carbohydrate moiety and the biosynthetic pathway towards cellulose synthesis. We show that the oxygen isotopic fractionation of the oxygen attached to carbon 2 of the glucose moieties differs from the average fractionation of the oxygens attached to carbons 3–6 from cellulose by at least 9%, for cellulose synthesized within seedlings of two different species (*Triticum aestivum* L. and *Ricinus communis* L.). The fractionation for a given oxygen in cellulose synthesized by the *Triticum* seedlings, which have starch as their primary carbon source, is different than the corresponding fractionation in *Ricinus* seedlings, within which lipids are the primary carbon source. This observation shows that the biosynthetic pathway towards cellulose affects oxygen isotope partitioning, a fact heretofore undemonstrated. Our findings may explain the species-dependent variability in the overall oxygen isotope fractionation during cellulose synthesis, and may provide much-needed insight for palaeoclimate reconstruction using fossil cellulose.

Key-words: oxygen-18; oxygen isotope ratios; palaeoclimate.

INTRODUCTION

Oxygen isotope ratios of plant cellulose are commonly invoked as a quantitative indicator of several abiotic and biotic factors integrated during cellulose synthesis. Initial isotopic studies of aquatic plant cellulose revealed that $\delta^{18}\text{O}$ values (see Eqn 2 for a definition of $\delta^{18}\text{O}$) of cellulose ($\delta^{18}\text{O}_{\text{cell}}$) were $\sim 27\%$ greater than the $\delta^{18}\text{O}$ value of water

(δ_{w}) at the site of synthesis (Epstein, Thompson & Yapp 1977; DeNiro & Epstein 1981). This has led to the conclusion that this enrichment factor results from a 2/3 contribution of ambient CO_2 (which has equilibrated with ambient water and has a $\delta^{18}\text{O}$ value $\sim 41\%$ greater than the $\delta^{18}\text{O}$ value of ambient water) and a 1/3 contribution from unfractionated environmental water (Epstein *et al.* 1977). Hence,

$$\delta^{18}\text{O}_{\text{cell}} = \left(\frac{1}{3} \cdot \delta_{\text{w}}\right) + \left[\frac{2}{3} \cdot (\delta_{\text{w}} + 41)\right] = \delta_{\text{w}} + 27.3\% \quad (1)$$

If Eqn 1 is valid, there should be a temperature effect on the isotopic composition of cellulose, because the isotopic equilibration between CO_2 and water is temperature sensitive (Vogel, Groates & Mark 1970). However, no such temperature effect has been observed, indicating that some additional processes have contributed to the oxygen isotopic signature of cellulose (DeNiro & Epstein 1981). Further evidence against a simplistic adoption of Eqn 1 is provided by tissue cultures, seed germination and other experiments within which cellulose synthesized heterotrophically showed post-photosynthetic isotopic exchange between the oxygen in intermediate carbohydrates and water (Sternberg, DeNiro & Savidge 1986; Yakir & DeNiro 1990; Luo & Sternberg 1992). Studies show that, depending on the substrate available for cellulose synthesis, 29–77% of the oxygen isotope content of heterotrophically synthesized cellulose is determined by the oxygen isotope content of water. Furthermore, cellulose synthesized from sucrose, the most common carbohydrate transported within plants, acquires about 40% of its oxygen from water (Sternberg *et al.* 1986; Yakir & DeNiro 1990; Roden & Ehleringer 1999). Current mechanistic models meant to explain $\delta^{18}\text{O}$ values in plant cellulose implicitly assume that the bulk of the isotopic signature acquired during cellulose synthesis occurs via the carbonyl hydration reaction. This assumption is supported by the observation that the fractionation factor for the carbonyl oxygen in an *in vitro* acetone–water system is 27‰ and is not subject to temperature effects (Sternberg & DeNiro 1983). The second implicit assumption is that oxygen isotope fractionation during heterotrophic cellulose synthesis relative to the *in situ* water remains constant regardless of the location of the oxygen in the carbohydrate moiety or the metabolic pathway leading to cellulose synthesis. Although widely used, these assumptions have not

Correspondence: Leonel Sternberg. Fax: +305 284 3039; e-mail: leo@bio.miami.edu

been tested, and several pathways other than the carbonyl hydration reaction by which the oxygen in the glucose moieties could acquire their particular isotopic signature during cellulose synthesis exists (Schmidt, Werner & Rossmann 2001). Furthermore, carbonyl hydration reactions may not achieve complete equilibrium (Schmidt *et al.* 2001). Some studies have indicated that the named oxygen isotopic fractionations vary and could be substantially below 27‰, as exemplified by the case of cellulose synthesized directly from sucrose and from starch (Sternberg *et al.* 1986; Luo & Sternberg 1992). These studies reported oxygen isotopic fractionation ranging from 16.3 to 22.0‰, as calculated for cellulose relative to the ^{18}O content of *in situ* water surrounding cellulose synthesis (Sternberg *et al.* 1986; Luo & Sternberg 1992). Although this evidence suggests that the second implicit assumption previously listed may not be valid, no study has yet been performed to explicitly test the true applicability of the assumption. Here, we report on experiments designed to specifically test the following assumptions: (1) the fractionation factors for oxygen atoms in a carbohydrate moiety are all similar to 27‰, regardless of the position of the oxygen within the carbohydrate molecule destined for cellulose synthesis, and (2) the metabolic pathway leading to cellulose synthesis is unimportant isotopically. We improved a previously developed technique (Sternberg, Anderson & Morrison 2003) so that we could determine the oxygen isotope ratios of the oxygen atoms specifically attached to the second carbon (*cf.* carbons 3–6) within the cellulose glucose moieties to test the first assumption. To test the second assumption, we compared the isotopic signature of cellulose that had been synthesized heterotrophically via two different biosynthetic pathways: using starch or lipid as the primary substrate. Using these approaches, we shed new light on the molecular and biosynthetic aspect of plant oxygen isotope fractionation.

METHODS AND MATERIALS

Heterotrophic cellulose synthesis and extraction

Cellulose was synthesized heterotrophically during the germination and growth of seedlings in complete darkness by two species: *Triticum aestivum* L. (within which starch comprises the major substrate for heterotrophic cellulose synthesis) and *Ricinus communis* L. (within which lipids comprise the major substrate for heterotrophic cellulose synthesis). We define heterotrophic cellulose synthesis as the synthesis of cellulose occurring in a sink tissue (e.g. stems, roots or seedlings grown in the dark) using carbohydrates transported from a source tissue (e.g. leaves or seed endosperm/cotyledons). Germination and growth of seedlings using water with variable oxygen isotopic ratios followed previous procedures (Sternberg *et al.* 1986; Luo & Sternberg 1992; Sternberg *et al.* 2003). After germination and stem/foilage growth, the fibrous stems, leaves and roots were separated, dried and ground to a fine powder. Cellulose extraction was performed with the classical methods described within Sternberg (1989a).

Cellulose hydrolysis and derivation

Extracted cellulose was hydrolyzed either with cellulase [enzyme class (EC 3.2.1.4)] or trifluoroacetic acid (TFA). For the enzymatic (cellulase) hydrolysis, we used a lower and more purified quantity of enzyme (Sigma-C-9422, 7.8 units mg^{-1} of solid, EC 3.2.1.4; Sigma-Aldrich, St. Louis, MO, USA) than previously reported (Sternberg *et al.* 2003). Although this product contains a high proportion of cellulase, it also contains cellobiase (EC 3.2.1.21) as per its activity specification (1 unit will liberate 1.0 μmol of glucose from cellulose in 1 h at pH 5.0 and 37 °C). One hundred milligrams of cellulase was added to 0.5 g of cellulose in 25 mL of a sodium acetate buffer (pH 5.0) and incubated at 40 °C for a period of 96 h. After incubation, the enzyme was precipitated by bringing the hydrolysis solution to a concentration of 75% MetOH, centrifuging and rotoevaporating the supernatant to recover the glucose. Acid hydrolysis was performed according to the method of Fengel & Wegener (1979) using 99% TFA, but with a smaller starting quantity of cellulose (0.3 g). Purity of the cellulose hydrolysate was confirmed by comparing its high-performance liquid chromatography (HPLC) chromatogram with that of a standard glucose solution via an Econosil-NH₂ 10 μ column (Alltech, Deerfield, IL, USA) with a 3:1 acetonitrile : water solution as the mobile phase. Sigma α -cellulose (C-8002, lot 11k0246; Sigma-Aldrich) was hydrolyzed by TFA using water of variable oxygen isotope signatures to determine whether the $\delta^{18}\text{O}$ of the hydrolysis solution affects the $\delta^{18}\text{O}$ value of the final phenylglucosazone derivative. Glucose from the cellulose hydrolysis was derived to phenylglucosazone using the method described within Oikawa *et al.* (1998).

Isotopic analysis of cellulose, phenylglucosazone and water

Oxygen isotope analyses of cellulose and phenylglucosazone were carried out as previously mentioned (Sternberg *et al.* 2003). Oxygen isotope ratios of the water available during seed germination and synthesis of heterotrophic cellulose were determined by equilibration of the water with CO₂ over a period of 12 h at 40 °C in a multiflow system connected to an Isoprime isotope ratio mass spectrometer (IRMS; GV Instruments, Manchester, UK). All oxygen isotope ratios reported here are in δ (‰) units, relative to Vienna Standard Mean Ocean Water (vSMOW) and are given by

$$\delta^{18}\text{O} = \left[\frac{R_{\text{sample}}}{R_{\text{vsmow}}} - 1 \right] \cdot 1000. \quad (2)$$

R_{sample} and R_{vsmow} represent the molar ratio of $^{18}\text{O}/^{16}\text{O}$ of the sample and Vienna-based standard mean ocean water. The precision of analysis was $\pm 0.36\text{‰}$ for the organic oxygen isotope determination (Sternberg *et al.* 2003) and $\pm 0.1\text{‰}$ for the water samples.

Data analysis

Calculation of the $\delta^{18}\text{O}$ value of the oxygen attached to carbon 2 of the glucose moieties of cellulose followed the procedure of Sternberg *et al.* (2003) and is described by the following equation:

$$\delta^{18}\text{O}^2 = (5 \times \delta^{18}\text{O}_{\text{cell}}) - (4 \times \delta^{18}\text{O}_{\text{p-g}}). \quad (3)$$

The terms of the previous equation: $\delta^{18}\text{O}^2$, $\delta^{18}\text{O}_{\text{cell}}$ and $\delta^{18}\text{O}_{\text{p-g}}$ represent the $\delta^{18}\text{O}$ values of the oxygen attached to the second carbon of the glucose moiety of cellulose, of cellulose and of phenylglucosazone, respectively. Regression values of slopes and intercepts for the linear relationship between $\delta^{18}\text{O}^2$, $\delta^{18}\text{O}_{\text{cell}}$ and $\delta^{18}\text{O}_{\text{p-g}}$ versus the $\delta^{18}\text{O}$ value of water available during seed germination and development were compared using the statistical methods of Armitage (1980). If the slopes were significantly different, then no comparison between the intercepts was performed. Error propagation for the calculations of the oxygen isotopic fractionation factor for each of the exchange reactions during cellulose synthesis was performed using the quadrature method (Taylor 1997).

RESULTS

Cellulose hydrolysis

A chromatogram of the TFA-generated cellulose hydrolysate was similar to that of the standard glucose solution, but with an additional minor peak appearing about 30 min after injection (Fig. 1). Oxygen isotope ratios of phenylglucosazone synthesized from the TFA-generated hydrolysates of Sigma α -cellulose averaged $23.4 \pm 2.2\%$, even though the hydrolysis solutions used spanned a range of $\delta^{18}\text{O}$ values over 250‰ (Fig. 2).

Oxygen isotope ratios of *Triticum* generated cellulose and phenylglucosazone

Oxygen isotope ratios of heterotrophically synthesized cellulose ($\delta^{18}\text{O}_{\text{cell}}$), phenylglucosazone ($\delta^{18}\text{O}_{\text{p-g}}$) and the




Figure 1. Ultraviolet (UV) absorption (292 nm) versus elution time for a standard glucose solution (heavy line) and cellulose trifluoroacetic acid (TFA) hydrolysate (light line).




Figure 2. Oxygen isotope ratios of phenylglucosazone prepared from Sigma α -cellulose hydrolysates versus the $\delta^{18}\text{O}$ values of the water within the hydrolysis solutions.

oxygen attached to the second carbon of the glucose moiety ($\delta^{18}\text{O}^2$) were all linearly related to the oxygen isotope ratios of the water (δ_w) available during seed germination and growth (Fig. 3, Table 1). There were, however, significant differences in the slopes of these linear relationships within the *Triticum* experiment. The linear relationship between $\delta^{18}\text{O}^2$ value versus the $\delta^{18}\text{O}$ value of water exhibited the highest slope (0.652 and 0.700 for the enzymatic and TFA hydrolysis, respectively) followed by that of cellulose (0.402) and finally, by that of phenylglucosazone (0.339 and 0.336 for the enzymatic and TFA hydrolysis, respectively). Although there were clear differences in the isotopic ratios of the y-intercept for the previously mentioned linear relationships (Table 1), they could not be compared statistically because of the extent to which the slope of the regressions differed.

A lower proportion of oxygen associated with carbon 2 exchanged with water during cellulose synthesis by the *Triticum* seedlings compared with a previous report (Sternberg *et al.* 2003). This discrepancy might be caused by the current use of a purer enzyme preparation, or by hydrolysis with TFA, which does not leave any residues. In Sternberg *et al.* (2003), a large quantity of low-purity enzyme (0.5 g) was used for hydrolysis and glucose, often present in some cellulase preparations, might have contaminated the hydrolysate. For these reasons, we prefer the values of isotopic exchange generated by this study as an improvement over Sternberg *et al.* (2003).

Oxygen isotope ratios of *Ricinus* generated cellulose and phenylglucosazone

No significant differences were found between the slopes for the linear relationships between $\delta^{18}\text{O}^2$, $\delta^{18}\text{O}_{\text{cell}}$ and $\delta^{18}\text{O}_{\text{p-g}}$ values versus the $\delta^{18}\text{O}$ value of the water available for the *Ricinus* generated cellulose (Fig. 3). A statistical comparison of the y-intercepts indicated significant differences: the $\delta^{18}\text{O}^2$ versus the $\delta^{18}\text{O}_w$ linear relationship had the highest y-intercept ($51.05 \pm 5.0\%$), followed by the intercept of the $\delta^{18}\text{O}_{\text{cell}}$ versus $\delta^{18}\text{O}_w$ relationship ($28.4 \pm 0.42\%$) and the




Figure 3. Oxygen isotope ratios of cellulose (a & b), phenylglucosazone (c & d) and the oxygen attached to carbon 2 of the glucose moieties (e & f) versus the $\delta^{18}\text{O}$ value of the water present during heterotrophic cellulose synthesis in both *Triticum* and *Ricinus* seedlings. Triangles (c & e) represent the case where cellulose was hydrolyzed enzymatically for the *Triticum* experiment; circles represent products from cellulose hydrolyzed by trifluoroacetic acid (TFA) for both experiments.

Table 1. The slope (F_i), intercept (I_i) and the correlation coefficient (R^2) for the relationship between the $\delta^{18}\text{O}$ value of cellulose, phenylglucosazone and oxygen attached to carbon 2 of the glucose moieties within heterotrophically synthesized cellulose versus that of the water within which the cellulose was synthesized for two species (*Triticum* and *Ricinus*)

Seed type	Oxygens in	F_i	I_i	R^2
<i>Triticum</i>	Cellulose	0.402 ± 0.007^a	28.6 ± 0.6	0.995
	Phenylglucosazone (enzyme)	0.339 ± 0.003^b	30.5 ± 0.2	0.999
	Phenylglucosazone (TFA)	0.336 ± 0.003^b	30.8 ± 0.2	0.999
	O in carbon 2 (enzyme)	0.652 ± 0.030^c	21.3 ± 3.2	0.944
	O in carbon 2 (TFA)	0.700 ± 0.030^c	19.6 ± 3.4	0.958
<i>Ricinus</i>	Cellulose	0.905 ± 0.005^a	28.4 ± 0.4^a	1.000
	Phenylglucosazone (TFA)	0.924 ± 0.012^a	22.5 ± 1.0^b	0.997
	O in carbon 2 (TFA)	0.858 ± 0.064^a	51.0 ± 5.0^c	0.948

In the case of *Triticum*, the cellulose was hydrolyzed either by enzymatic hydrolysis or by TFA (see Methods and Materials). The slope F_i represents the proportion of oxygen that was either exchanged or added by water during cellulose synthesis. Slope and/or intercept values superscripted for each species indicate a significant difference between slopes and/or intercept. In *Triticum*, the slopes were significantly different and therefore, the difference between intercepts could not be tested. TFA, trifluoroacetic acid.

lowest intercept for the phenylglucosazone derivative ($22.5 \pm 1.0\%$).

DISCUSSION

Cellulose hydrolyses by enzyme and TFA

Cellulose synthesized from *Triticum* seeds were hydrolyzed in two ways: with an enzyme preparation or with TFA. We observed no significant differences between the oxygen isotope ratios of phenylglucosazone ($\delta^{18}\text{O}_{\text{P-g}}$), and the oxygen attached to carbon 2 ($\delta^{18}\text{O}^2$) for cellulose hydrolyzed by either method. Therefore, even though the enzyme incompletely hydrolyzed the cellulose, there are no fractionation effects from the incomplete hydrolysis.

The cellulose hydrolysis by TFA yielded a hydrolysate highly concentrated in glucose with a minor component (probably mannose) eluting approximately 5 min after the glucose eluted (Fig. 1). Cellulose hydrolysis by TFA is a much faster method of hydrolysis compared with the enzymatic method: cellulose can be completely hydrolyzed in 2 h with TFA, whereas enzymatic hydrolysis requires an incubation period of nearly 4 d. In addition, the TFA method yields a much greater quantity of hydrolysate than the enzymatic method and requires less starting material, which can be a critical consideration when analyzing fossil samples (Jahren & Sternberg 2003). Lastly, cellulose hydrolysis by TFA ensures against contamination of the

hydrolysate, because TFA is completely removed during rotoevaporation (Fengel & Wegener 1979).

Isotopic effects during cellulose hydrolysis

It is important to determine any isotopic signal originating from the hydrolysis of cellulose when considering the phenylglucosazone derivative. If such a signal exists, it could complicate the use of the isotopic ratios of phenylglucosazone to interpret isotopic values resulting from cellulose synthesis within plants. Cellulose hydrolysis involves the insertion of one water molecule between two consecutive glucose moieties (Fig. 4). A hydroxyl group, which contains one oxygen atom, could be added either to carbon 1 or 4 of the glucose moieties (Fig. 4). Hydrolysis of cellulose with ^{18}O -enriched water, followed by the derivation and isotopic analysis of phenylglucosazone, can determine which carbon of the glucose moiety receives an oxygen atom from water. If the ^{18}O from the hydrolysis solution is added to carbon 1, no isotopic enrichment will be apparent because of the fact that both oxygens attached to carbon 1 and 2 are eliminated during derivation of phenylglucosazone (Fig. 4). In contrast, if the oxygen atom from water is being added to the carbon 4 of the glucose moiety, phenylglucosazone will have a greater $\delta^{18}\text{O}$ value than that produced from hydrolysis of cellulose in ^{18}O -depleted water (Fig. 4). No significant difference between the $\delta^{18}\text{O}$ values of phenylglucosazone derived from cellulose hydrolysates in the




Figure 4. Comparison of possible outcomes from the addition of a hydroxyl group to two different positions (carbons 1 or 4) of the glucose moieties during the hydrolysis of cellulose.

presence of water having oxygen isotope ratios spanning 250‰ was observed (Fig. 2). From this, we conclude that the hydroxyl group from the hydrolysis solution is added to carbon 1 of the glucose moiety and was then eliminated when phenylglucosazone was derived. This means that the phenylglucosazone isotopic signal results exclusively from *in situ* biochemical reactions that occur during cellulose synthesis in plants and not from other processes that take place during hydrolysis.

Relationship between $\delta^{18}\text{O}$ values of cellulose and intramolecular oxygen with those of germination water

The linear regression of $\delta^{18}\text{O}_{\text{cell}}$, $\delta^{18}\text{O}_{\text{p-g}}$ and $\delta^{18}\text{O}^2$ values against the $\delta^{18}\text{O}$ value of water present during germination and seedling growth (δ_w) can be understood as the following (Sternberg *et al.* 1986):

$$\delta^{18}\text{O}_{\text{cell}} = [F_{\text{cell}} \cdot \delta_w] + [(F_{\text{cell}} \cdot \Delta_{\text{cell}}) + (1 - F_{\text{cell}}) \cdot \delta_{\text{cell}}^{\text{nc}}] \quad (4)$$

$$\delta^{18}\text{O}_{\text{p-g}} = [F_{\text{p-g}} \cdot \delta_w] + [(F_{\text{p-g}} \cdot \Delta_{\text{p-g}}) + (1 - F_{\text{p-g}}) \cdot \delta_{\text{p-g}}^{\text{nc}}] \quad (5)$$

$$\delta^{18}\text{O}^2 = [F_2 \cdot \delta_w] + [(F_2 \cdot \Delta_2) + (1 - F_2) \cdot \delta_2^{\text{nc}}] \quad (6)$$

The slope of the linear relationship (F_i) is equivalent to the proportion of oxygen in the substrate (i.e. starch or lipid) that either exchanged with or were added by water during cellulose synthesis. These values are specific to the oxygen atoms in the cellulose molecule (attached to carbons 2–6 for F_{cell} , carbons 3–6 for $F_{\text{p-g}}$ and carbon 2 for F_2). The overall isotopic fractionation for the oxygen that exchanged with or was added by water during cellulose synthesis (Δ_{cell}) is composed of a 4/5 contribution from the overall fractionation of the oxygen associated with carbons 3–6 ($\Delta_{\text{p-g}}$) and a 1/5 contribution from the fractionation in the oxygen associated with carbon 2 (Δ_2). Likewise, the isotopic value of the oxygen atoms in the substrate that does not exchange with or are not added by water is designated as δ_i^{nc} for the respective oxygen atoms in the glucose moieties. The y-intercept of the previous linear relationships (designated here as I_i) is equivalent to the isotopic fraction of the oxygen that did not exchange with, or was not added on, by water during cellulose synthesis $[(1 - F_i) \cdot \delta_i^{\text{nc}}]$, as well as to a fractionation factor component ($F_i \Delta_i$).

Proportion of intramolecular oxygen exchanging with water during heterotrophic cellulose synthesis

We observed a greater proportion of isotopic exchange between the oxygen attached to carbon 2 and water compared with the proportional exchange of oxygen attached to carbons 3–6 during cellulose synthesis by *Triticum*. The isotopic exchange during cellulose synthesis is expected to be greater in the oxygen associated with carbon 2 than with those attached to carbons 3–6, because glucose molecules from starch would have to first be converted to fructose-bis-phosphate, thus exposing the oxygen attached to carbon 2 to exchange, before it is converted to a triose phosphate

and exposing the oxygen attached to carbons 3–6 to exchange.

Our results imply that complete isotopic exchange occurred between the oxygen in the *Ricinus* (i.e. oxygen-poor lipid substrate) synthesized cellulose and water. The slope of the linear relationship of cellulose and derivatives against the *in situ* water ranged from 0.86 to 0.92. Previous experiments using *Ricinus* seeds resulted in a lower slope of the linear relationship, indicating that only about 70% of the cellulose oxygen molecules had been isotopically labeled by water during its synthesis (Sternberg *et al.* 1986; Luo & Sternberg 1992). Oxygen isotope analyses in both of these previous experiments were performed using isoquinoline (Sternberg 1989a), which carries an oxygen isotopic 'memory effect', thus dampening isotopic labelling and lowering the slope of the relationship relative to that presented here. There are no reports describing a 1:1 relationship between the $\delta^{18}\text{O}$ value of autotrophically synthesized cellulose and that of water. Autotrophically synthesized cellulose should have its oxygen isotope ratios completely determined by the oxygen isotope ratios of the *in situ* water. Previous studies of autotrophically synthesized cellulose usually measured slopes between 0.48 and 0.88 (Cooper & DeNiro 1989; Yakir & DeNiro 1990; Sauer, Miller & Overpeck 2001). Previous explanations proposed that a change in the isotopic composition of the *in situ* water available during the growth of the plant material caused the slope to be less than expected (Yakir & DeNiro 1990). However, a different autotrophic cellulose experiment showed that the changes in the $\delta^{18}\text{O}$ value of water during plant growth were too small to account for a slope of less than 1 (Sauer *et al.* 2001). Our results show that all the oxygen atoms in the heterotrophically synthesized cellulose in the *Ricinus* experiment were derived from the *in situ* water, because the slopes we obtained were similar to, or higher than, those observed for experiments that generated cellulose autotrophically.

Intramolecular and biosynthetic effects in the oxygen isotope fractionation during heterotrophic cellulose synthesis

We tested the assumption that the oxygen isotopic fractionation between oxygen in the carbohydrate intermediates and oxygen within the *in situ* water is approximately 27‰ regardless of (1) the oxygen position in the carbohydrate molecule and (2) the metabolic pathway leading to cellulose. If the first assumption was correct, then the y-intercepts of Eqns 5 and 6 can be written as

$$I_{\text{p-g}} = (F_{\text{p-g}} \cdot \Delta) + (1 - F_{\text{p-g}}) \cdot \delta_{\text{p-g}}^{\text{nc}} \quad \text{and} \quad (7)$$

$$I_2 = (F_2 \cdot \Delta) + (1 - F_2) \cdot \delta_2^{\text{nc}}, \quad \text{respectively.} \quad (8)$$

For Eqns 7 and 8 above, Δ represents a single fractionation factor regardless of the oxygen position in the carbohydrate leading to cellulose synthesis (Case 1). In the same vein, assuming that the non-exchangeable oxygen attached to

carbon 2 and those attached to carbons 3–6 are similar to the respective oxygen of the original substrate (e.g. starch), it follows that

$$\delta_{\text{starch}} = (0.20 \cdot \delta_2^{\text{nc}}) + (0.80 \cdot \delta_{\text{p-g}}^{\text{nc}}). \quad (9)$$

We measured the $\delta^{18}\text{O}$ value of starch from the *Triticum* seedlings (δ_{starch}). Therefore, Eqns 7–9 only have three unknowns (Δ , δ_2^{nc} and $\delta_{\text{p-g}}^{\text{nc}}$) and can be solved simultaneously for Δ . Calculation of Δ under these assumptions showed a mean fractionation factor of 23.5‰ (Table 2), which is low compared to the 27‰ observed during autotrophic cellulose synthesis (Epstein *et al.* 1977; DeNiro & Epstein 1981). A small error is associated with Eqn 9, because about 70% of starch is composed of amylopectin with branch lengths of about 20 glucose moieties (Blennow *et al.* 2000). This shorter branch length would be associated with several oxygen atoms attached to the first carbon of the terminal glucose moieties. As a result, the factors of 0.20 and 0.80 previously mentioned would be an overestimate by a small amount, and the oxygen isotopic signature of the oxygen attached to carbon 1 would be unknown. However, a sensitivity analysis, assuming that the $\delta^{18}\text{O}$ value of the oxygen atoms in carbon 1 of the glucose moieties deviated from the average of starch by 100‰, gave an error of only $\sim \pm 0.09\%$.

The assumption that the fractionation factors between each oxygen in the glucose moieties of cellulose are similar, regardless of the oxygen position, leads to a contradiction of previous well-established observations and is therefore shown to be false. Further evidence that the fractionation factor for the oxygen associated with carbon 2 is different than the average for the oxygen associated with carbons 3–6 is provided when these two fractionation factors are calculated independently (Case 2). The only required condition for this calculation is that the relative contribution of each fractionation factor will add up to the reported average oxygen isotopic fractionation of autotrophically synthesized cellulose: 27‰ (DeNiro & Epstein 1981; Sternberg 1989b). Hence,

$$\Delta_{\text{cell}} = 27\text{‰} = (0.20 \cdot \Delta_2) + (0.80 \cdot \Delta_{\text{p-g}}) \quad (10)$$

and the y-intercept in equations 5 and 6 become

$$I_2 = (F_2 \cdot \Delta_2) + (1 - F_2) \cdot \delta_2^{\text{nc}} \quad \text{and} \quad (11)$$

$$I_{\text{p-g}} = (F_{\text{p-g}} \cdot \Delta_{\text{p-g}}) + (1 - F_{\text{p-g}}) \cdot \delta_{\text{p-g}}^{\text{nc}} \quad (12)$$

Equations 9–12 have four unknowns: the fractionation factors Δ_2 and $\Delta_{\text{p-g}}$, and the non-exchangeable oxygen isotope ratios of the substrate oxygen associated with carbon 2 (δ_2^{nc}) and 3–6 ($\delta_{\text{p-g}}^{\text{nc}}$). These four equations can be solved simultaneously for the unknowns, including Δ_2 and $\Delta_{\text{p-g}}$. The resulting fractionation factor for the oxygen associated with carbon 2 (averaging 19.6‰) is different than the fractionation for the oxygen associated with carbons 3–6 (averaging 28.8‰, Table 2), therefore negating the first assumption.

The results of the *Ricinus* experiment, where lipids are the major substrate for cellulose synthesis, provide evidence that fractionation factors not only differ within the cellulose molecule, but also depend on the metabolic pathway by which cellulose is synthesized. This possibility was first mentioned by Schmidt *et al.* (2001); here we quantify this effect. The oxygen isotope fractionation factors resulting from the *Ricinus* experiment show large differences between oxygen associated with carbon 2 compared with the oxygen associated with carbons 3–6 of the glucose moieties (Table 2). The metabolic pathway from lipid to cellulose in *Ricinus* is obviously different than the pathway transforming starch into cellulose in *Triticum*. The observation that the fractionation for the oxygen associated with carbon 2 in the *Ricinus* experiment ($\Delta_2 = 51\%$) differed greatly from that observed for the *Triticum* experiment ($\Delta_2 = 19.6\%$) leads us to the conclusion that the particular cellulose biosynthetic pathway exerts an influence over the fractionation effects of individual oxygen atoms, therefore negating the second assumption previously detailed. The differences between cellulose synthesized from starch versus that synthesized from lipids in terms of oxygen addition or exchange lies in several reactions occurring during

Table 2. Fractionation factors for the oxygen attached to carbon 2 (Δ_2), the average fractionation for oxygen attached to carbons 3–6 ($\Delta_{\text{p-g}}$) and the average fractionation factor for all the oxygen within a cellulose molecule (Δ_{cell}), as well as the oxygen isotope ratios of the different oxygen atoms in the starch substrate for the *Triticum* experiment

Seed type and hydrolysis method	Δ_2	$\Delta_{\text{p-g}}$	Δ_{cell}	Starch δ_2	Starch $\delta_{\text{p-g}}$
<i>Triticum</i>					
Case 1					
Enzyme hydrolysis	24.0 ± 2.9	24.0 ± 2.9	24.0 ± 2.9	16.4	33.7
TFA hydrolysis	22.9 ± 3.1	22.9 ± 3.1	22.9 ± 3.1	12.0	34.8
Case 2					
Enzyme hydrolysis	18.9 ± 2.6	29.0 ± 3.9	27.0	26.5	31.2
TFA hydrolysis	20.2 ± 2.7	28.7 ± 3.9	27.0	23.8	31.9
<i>Ricinus</i>					
TFA hydrolysis	51.0 ± 5.0	22.5 ± 1.0	28.2 ± 0.4	n.a.	n.a.

For the *Triticum* experiment, fractionation factors were calculated for two cases: case 1 assumed that all fractionation factors are the same regardless of the position in the glucose moiety, and case 2 made no assumption on the fractionation factors other than they must average to 27‰. For the *Ricinus* experiment, we assumed complete addition/exchange between the oxygen of the glucose moieties and water during cellulose synthesis, that is, $F \approx 1$.

n.a., not applicable; TFA, trifluoroacetic acid.

gluconeogenesis from lipids that do not occur during the biosynthetic pathway of cellulose synthesis from starch. Glucose moieties generated from starch hydrolysis have all the oxygen atoms found in the final cellulose product, and their final isotopic identity is mostly acquired during carbonyl hydration and the aldolase catalyzed reaction occurring during the futile cycle from glucose-6-phosphate to the triose phosphates (Schmidt *et al.* 2001). Carbonyl hydration reactions may be facilitated and may reach equilibrium through enzyme catalysis (Schmidt *et al.* 2001). Glucose moieties generated from lipids, however, will have all the oxygen atoms derived from water through reactions catalyzed by enolhydratase, fumarase and enolase, and the final isotopic identity of the oxygen atoms in the cellulose molecule may be determined by reactions during which the newly added oxygen is incompletely exchanged with water (Schmidt *et al.* 2001). It has been suggested, for example, that the oxygen added during gluconeogenesis in position 6 of the glucose moiety does not exchange with water, because it is protected by a phosphate group (Schmidt *et al.* 2001).

The source of variation in oxygen isotope fractionation during cellulose synthesis

The commonly used average 27‰ enrichment factor for the oxygen isotope ratios of cellulose relative to the source water is based on several measurements of the oxygen isotope ratios of cellulose and the source water of submerged aquatic plants (DeNiro & Epstein 1981; Sternberg, DeNiro & Keeley 1984; Sternberg 1989b; Sauer *et al.* 2001). This value is, in fact, a mean of the previous measurements having a range of about $\pm 4\%$ (DeNiro & Epstein 1981; Sternberg 1989b). The reasons for this variation is as yet unknown, but careful monitoring of source water indicates that this variation is not due to isotopic drift of the source water during the growth of the aquatic plant (Sauer *et al.* 2001). Furthermore, different species grown within the same aquarium will have different cellulose oxygen isotope ratios (Sternberg *et al.* 1984). We propose that the intramolecular and the biosynthetic pathway isotopic effects demonstrated here may in part be responsible for the previously observed variation. The differences in the fractionations associated with particular oxygen positions in the glucose moiety can affect the final average cellulose oxygen isotope ratio relative to that of the source water. There are likely differences in the extent of the (hexose-triose) futile cycle (Hill *et al.* 1995) occurring in the cellulose synthesizing tissue, depending on species and even environmental conditions (Saurer, Aellen & Siegwolf 1997). This difference would not only affect the proportion of oxygen atoms in the glucose moiety that interact with the water in the cellulose synthesizing cell, but it would also affect which oxygen atom interacts with the cell water. The net outcome will be a difference in the average isotopic fractionation during cellulose synthesis, because the isotopic fractionations for different oxygen positions in the glucose differ. Biosynthetic pathways are also important in determining the final

oxygen isotope ratios of cellulose even if the immediate precursor for cellulose synthesis is a carbohydrate (e.g. sucrose). We note that, although the ultimate carbon source for cellulose synthesis in the *Ricinus* seedlings is lipids, the pathway towards cellulose synthesis probably involves gluconeogenesis from lipids stored in the cotyledons, conversion to sucrose, translocation to sink tissue and finally, cellulose synthesis. Evidently, the isotopic labelling occurring during gluconeogenesis is sufficiently robust to persist through the entire cellulose synthesis process, including the sucrose intermediate, and to affect the oxygen isotope ratios of cellulose: the final biosynthetic product.

ACKNOWLEDGMENTS

This research was supported in part by NSF grant OPP/0323877 to A.H. Jahren and NSF grant OPP/0322364 to L.S.L. Sternberg. We thank W.M. Hagopian for his laboratory expertise.

REFERENCES

- Armitage P. (1980) *Statistical Methods in Medical Research*, pp. 279–301. Blackwell Scientific Publications, Oxford UK.
- Blennow A., Engelsen S.B., Munck L. & Møller B.L. (2000) Starch molecular structure and phosphorylation investigated by a combined chromatographic and chemometric approach. *Carbohydrate Polymers* **41**, 163–174.
- Cooper L.W. & DeNiro M.J. (1989) Oxygen-18 content of atmospheric oxygen does not affect the oxygen isotope relationship between environmental water and cellulose in a submerged aquatic plant, *Egeria densa* Planch. *Plant Physiology* **91**, 536–541.
- DeNiro M.J. & Epstein S. (1981) Isotopic composition of cellulose from aquatic organisms. *Geochimica et Cosmochimica Acta* **45**, 1894–1981.
- Epstein S., Thompson P. & Yapp C.J. (1977) Oxygen and hydrogen isotopic ratios in plant cellulose. *Science* **198**, 1209–1215.
- Fengel D. & Wegener G. (1979) Hydrolysis of polysaccharides with trifluoroacetic acid and its application to rapid wood and pulp analysis. In *Hydrolysis of Cellulose; Mechanism of Enzymatic and Acid Catalysis* (eds R.D. Brown Jr. & L. Jurassek) Advances in Chemistry Series No 181, pp. 145–158). American Chemical Society.
- Hill S.A., Waterhouse J.S., Field E.M., Switsur V.R. & ap Rees T. (1995) Rapid recycling of triose phosphates in oak stem tissue. *Plant, Cell & Environment* **18**, 931–936.
- Jahren A.H. & Sternberg L.S.L. (2003) Humidity estimate for the middle-Eocene Arctic rainforest. *Geology* **31**, 463–466.
- Luo Y.H. & Sternberg L.S.L. (1992) Hydrogen and oxygen isotopic fractionation during heterotrophic cellulose synthesis. *Journal of Experimental Botany* **43**, 47–50.
- Oikawa N., Muller C., Kunz M. & Lichtenthaler F.W. (1998) Hydrophically functionalized pyrazoles from sugars. *Carbohydrate Research* **309**, 269–279.
- Roden J.S. & Ehleringer J.R. (1999) Hydrogen and oxygen isotope ratios of tree-ring cellulose for riparian trees grown long-term under hydroponically controlled environments. *Oecologia* **121**, 467–477.
- Sauer P.E., Miller G.M. & Overpeck J.T. (2001) Oxygen isotope ratios of organic matter in arctic lakes as a paleoclimate proxy: field and laboratory investigations. *Journal of Paleolimnology* **25**, 43–64.

- Saurer M., Aellen K. & Siegwolf R. (1997) Correlating $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in cellulose of trees. *Plant, Cell & Environment* **20**, 1543–1550.
- Schmidt H.-L., Werner R.A. & Rossmann A. (2001) ^{18}O pattern and biosynthesis of natural plant products. *Phytochemistry* **58**, 9–32.
- Sternberg L.S.L. (1989a) Oxygen and hydrogen isotope measurements in plant cellulose analysis. In *Modern Methods of Plant Analysis*, New series Vol. 10, Plant fibres (eds H.F. Linskens & J.F. Jackson) pp. 89–98. Springer Verlag, NY, USA.
- Sternberg L.S.L. (1989b) Oxygen and hydrogen isotope ratios in plant cellulose: mechanisms and applications. In *Stable isotope in ecological research, Ecological studies 68* (eds P.W. Rundel, J.R. Ehleringer & K.A. Nagy), pp. 124–141. Springer Verlag, NY, USA.
- Sternberg L., Da S.L. & DeNiro M.J. (1983) Biogeochemical implications of the isotopic equilibrium fractionation factor between oxygen atoms of acetone and water. *Geochimica Cosmochimica Acta* **47**, 2271–2274.
- Sternberg L.S.L., DeNiro M.J. & Keeley J.E. (1984) Hydrogen, oxygen and carbon isotope ratios of cellulose from submerged aquatic crassulacean acid metabolism and non-crassulacean acid metabolism plants. *Plant Physiology* **76**, 68–70.
- Sternberg L.S.L., DeNiro M.J. & Savidge R.A. (1986) Oxygen isotope exchange between metabolites and water during biochemical reactions leading to cellulose synthesis. *Plant Physiology* **82**, 423–427.
- Sternberg L.S.L., Anderson W.T. & Morrison K. (2003) Separating soil and leaf water ^{18}O isotopic signals in plant stem cellulose. *Geochimica Cosmochimica Acta* **67**, 2561–2566.
- Taylor R.J. (1997) *An Introduction to Error Analysis*, p. 327. University Science Books, Sausalito, CA.
- Vogel J.C., Groates P.M. & Mark W.G. (1970) Isotopic fractionations between gaseous and dissolved carbon dioxide. *Zeitschrift für Physik* **230**, 225–238.
- Yakir D. & DeNiro M.J. (1990) Oxygen and hydrogen isotopic fractionation during cellulose metabolism in *Lemna gibba* L. *Plant Physiology* **93**, 325–332.

Received 26 February 2006; received in revised form 20 April 2006;
Accepted for publication 3 May 2006