

American Geophysical Union Fall Meeting, San Francisco, CA, Dec 12-16, 2016

Combined and isolated effects of $p\text{CO}_2$ and soil water content on carbon isotope discrimination during C_3 photosynthesis

Brian A. Schubert¹ and A. Hope Jahren²

¹*School of Geosciences, University of Louisiana at Lafayette, Lafayette, LA 70504;*

²*Centre for Earth Evolution and Dynamics, University of Oslo, N-0315, Oslo, Norway*

Biomass produced via C_3 photosynthesis dominates the terrestrial organic matter (TOM) found within the geologic record. Our previous work revealed an increase in net discrimination ($\Delta^{13}\text{C}$) $\approx +4\text{‰}$ across an increase in $p\text{CO}_2$ level from ambient to $\text{RCO}_2 = 6\text{x}$ within the model C_3 plant *Arabidopsis thaliana*, grown to maturity under constant conditions of light, moisture, and nutrient availability (Schubert and Jahren, 2012, *GCA*), leading us to suggest that changes in ancient $p\text{CO}_2$ level can be reconstructed from $\Delta^{13}\text{C}$ within terrestrial sediments. Others have observed an average change in $\Delta^{13}\text{C} \approx +4\text{‰}$ when comparing the $\delta^{13}\text{C}$ value of herbarium samples collected from cool-cold forests to tropical environments against the MAP recorded (Diefendorf et al., 2010, *PNAS*), leading those authors to suggest that changes in the $\Delta^{13}\text{C}$ value of TOM recovered from the geological record can be interpreted as changes in precipitation level and/or water availability.

Because decreasing moisture availability and increasing $p\text{CO}_2$ level exert control over $\Delta^{13}\text{C}$ through distinctly different mechanisms (i.e., decreased stomatal conductance vs. inhibition of photorespiration, respectively), a simultaneous change in both $p\text{CO}_2$ level and moisture availability could combine to influence carbon isotope fractionation. Here we present experiments in which we grew 230 *A. thaliana* plants at each of 5 levels of $p\text{CO}_2$: 390, 685, 1075, 1585, and 2175 ppmv. Within each growth chamber, soil moisture content (θ_m) was maintained at 1.50 g g^{-1} for 11 days following germination. Afterwards, we allowed 170 of the plants to dry to $\theta_m = 0.83, 0.44,$ and 0.38 g g^{-1} . After 3 weeks of total growth, tissues were analyzed for $\delta^{13}\text{C}$ value. We compare the isolated and combined effects of $p\text{CO}_2$ and soil moisture upon carbon isotope fractionation across the total range of $p\text{CO}_2$ levels reconstructed for the last 350 million years and across moisture levels associated with a $\sim 4.5\text{x}$ change in plant biomass.